

JMAP REGENTS BY TYPE

The NY Algebra 2/Trigonometry Regents Exams
Fall 2009-June 2014

www.jmap.org

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

1 For which value of k will the roots of the equation $2x^2 - 5x + k = 0$ be real and rational numbers?

- 1) 1
- 2) -5
- 3) 0
- 4) 4

2 What is the range of $f(x) = |x - 3| + 2$?

- 1) $\{x|x \geq 3\}$
- 2) $\{y|y \geq 2\}$
- 3) $\{x|x \in \text{real numbers}\}$
- 4) $\{y|y \in \text{real numbers}\}$


3 Which expression represents the total number of different 11-letter arrangements that can be made using the letters in the word "MATHEMATICS"?

- 1) $\frac{11!}{3!}$
- 2) $\frac{11!}{2! + 2! + 2!}$
- 3) $\frac{11!}{8!}$
- 4) $\frac{11!}{2! \cdot 2! \cdot 2!}$


4 In parallelogram $BFLO$, $OL = 3.8$, $LF = 7.4$, and $m\angle O = 126$. If diagonal BL is drawn, what is the area of $\triangle BLF$?

- 1) 11.4
- 2) 14.1
- 3) 22.7
- 4) 28.1


5 If $m\angle\theta = -50$, which diagram represents θ drawn in standard position?




1)



2)



3)




4)

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org


- 6 The expression $\sin(\theta + 90)^\circ$ is equivalent to
- 1) $-\sin \theta$
 - 2) $-\cos \theta$
 - 3) $\sin \theta$
 - 4) $\cos \theta$
- 7 What is the equation of the circle passing through the point $(6, 5)$ and centered at $(3, -4)$?
- 1) $(x - 6)^2 + (y - 5)^2 = 82$
 - 2) $(x - 6)^2 + (y - 5)^2 = 90$
 - 3) $(x - 3)^2 + (y + 4)^2 = 82$
 - 4) $(x - 3)^2 + (y + 4)^2 = 90$
- 8 If n is a negative integer, then which statement is always true?
- 1) $6n^{-2} < 4n^{-1}$
 - 2) $\frac{n}{4} > -6n^{-1}$
 - 3) $6n^{-1} < 4n^{-1}$
 - 4) $4n^{-1} > (6n)^{-1}$
- 9 The discriminant of a quadratic equation is 24. The roots are
- 1) imaginary
 - 2) real, rational, and equal
 - 3) real, rational, and unequal
 - 4) real, irrational, and unequal
- 10 Max solves a quadratic equation by completing the square. He shows a correct step:
- $$(x + 2)^2 = -9$$
- What are the solutions to his equation?
- 1) $2 \pm 3i$
 - 2) $-2 \pm 3i$
 - 3) $3 \pm 2i$
 - 4) $-3 \pm 2i$
- 11 Liz has applied to a college that requires students to score in the top 6.7% on the mathematics portion of an aptitude test. The scores on the test are approximately normally distributed with a mean score of 576 and a standard deviation of 104. What is the minimum score Liz must earn to meet this requirement?
- 1) 680
 - 2) 732
 - 3) 740
 - 4) 784
- 12 A cliff diver on a Caribbean island jumps from a height of 105 feet, with an initial upward velocity of 5 feet per second. An equation that models the height, $h(t)$, above the water, in feet, of the diver in time elapsed, t , in seconds, is
- $$h(t) = -16t^2 + 5t + 105.$$
- How many seconds, to the *nearest hundredth*, does it take the diver to fall 45 feet below his starting point?
- 1) 1.45
 - 2) 1.84
 - 3) 2.10
 - 4) 2.72

- 13 What is the equation of the graph shown below?



- 1) $y = 2^x$
- 2) $y = 2^{-x}$
- 3) $x = 2^y$
- 4) $x = 2^{-y}$

- 14 Which equation represents the graph below?



- 1) $y = -2 \sin 2x$
- 2) $y = -2 \sin \frac{1}{2}x$
- 3) $y = -2 \cos 2x$
- 4) $y = -2 \cos \frac{1}{2}x$

- 15 The roots of the equation $2x^2 + 4 = 9x$ are
- 1) real, rational, and equal
 - 2) real, rational, and unequal
 - 3) real, irrational, and unequal
 - 4) imaginary

- 16 When factored completely, $x^3 + 3x^2 - 4x - 12$ equals
- 1) $(x + 2)(x - 2)(x - 3)$
 - 2) $(x + 2)(x - 2)(x + 3)$
 - 3) $(x^2 - 4)(x + 3)$
 - 4) $(x^2 - 4)(x - 3)$

- 17 A market research firm needs to collect data on viewer preferences for local news programming in Buffalo. Which method of data collection is most appropriate?
- 1) census
 - 2) survey
 - 3) observation
 - 4) controlled experiment

- 18 What is the number of degrees in an angle whose radian measure is $\frac{8\pi}{5}$?
- 1) 576
 - 2) 288
 - 3) 225
 - 4) 113

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

19 A sequence has the following terms: $a_1 = 4$, $a_2 = 10$, $a_3 = 25$, $a_4 = 62.5$. Which formula represents the n th term in the sequence?

- 1) $a_n = 4 + 2.5n$
- 2) $a_n = 4 + 2.5(n - 1)$
- 3) $a_n = 4(2.5)^n$
- 4) $a_n = 4(2.5)^{n-1}$

20 The expression $\frac{a + \frac{b}{c}}{d - \frac{b}{c}}$ is equivalent to

- 1) $\frac{c+1}{d-1}$
- 2) $\frac{a+b}{d-b}$
- 3) $\frac{ac+b}{cd-b}$
- 4) $\frac{ac+1}{cd-1}$

21 A circle is drawn to represent a pizza with a 12 inch diameter. The circle is cut into eight congruent pieces. What is the length of the outer edge of any one piece of this circle?

- 1) $\frac{3\pi}{4}$
- 2) π
- 3) $\frac{3\pi}{2}$
- 4) 3π

22 What is the solution set for the equation

$$\sqrt{5x+29} = x+3?$$

- 1) $\{4\}$
- 2) $\{-5\}$
- 3) $\{4, 5\}$
- 4) $\{-5, 4\}$

23 A school math team consists of three juniors and five seniors. How many different groups can be formed that consist of one junior and two seniors?

- 1) 13
- 2) 15
- 3) 30
- 4) 60

24 If $\log x^2 - \log 2a = \log 3a$, then $\log x$ expressed in terms of $\log a$ is equivalent to

- 1) $\frac{1}{2} \log 5a$
- 2) $\frac{1}{2} \log 6 + \log a$
- 3) $\log 6 + \log a$
- 4) $\log 6 + 2 \log a$

25 Expressed in simplest form, $\frac{3y}{2y-6} + \frac{9}{6-2y}$ is equivalent to

- 1) $\frac{-6y^2 + 36y - 54}{(2y-6)(6-2y)}$
- 2) $\frac{3y-9}{2y-6}$
- 3) $\frac{3}{2}$
- 4) $-\frac{3}{2}$

26 A jogger ran $\frac{1}{3}$ mile on day 1, and $\frac{2}{3}$ mile on day 2, and $1\frac{1}{3}$ miles on day 3, and $2\frac{2}{3}$ miles on day 4, and this pattern continued for 3 more days. Which expression represents the total distance the jogger ran?

- 1) $\sum_{d=1}^7 \frac{1}{3} (2)^{d-1}$
- 2) $\sum_{d=1}^7 \frac{1}{3} (2)^d$
- 3) $\sum_{d=1}^7 2 \left(\frac{1}{3}\right)^{d-1}$
- 4) $\sum_{d=1}^7 2 \left(\frac{1}{3}\right)^d$


27 A school cafeteria has five different lunch periods. The cafeteria staff wants to find out which items on the menu are most popular, so they give every student in the first lunch period a list of questions to answer in order to collect data to represent the school. Which type of study does this represent?

- 1) observation
- 2) controlled experiment
- 3) population survey
- 4) sample survey


28 The expression $(2a)^{-4}$ is equivalent to

- 1) $-8a^4$
- 2) $\frac{16}{a^4}$
- 3) $-\frac{2}{a^4}$
- 4) $\frac{1}{16a^4}$


29 Which is a graph of $y = \cot x$?




1)



2)



3)



4)

30 What is the solution set of the equation

$$-\sqrt{2} \sec x = 2 \text{ when } 0^\circ \leq x < 360^\circ?$$

- 1) $\{45^\circ, 135^\circ, 225^\circ, 315^\circ\}$
- 2) $\{45^\circ, 315^\circ\}$
- 3) $\{135^\circ, 225^\circ\}$
- 4) $\{225^\circ, 315^\circ\}$

31 If $g(x) = \frac{1}{2}x + 8$ and $h(x) = \frac{1}{2}x - 2$, what is the value of $g(h(-8))$?

- 1) 0
- 2) 9
- 3) 5
- 4) 4

32 The expression $\log 4m^2$ is equivalent to

- 1) $2(\log 4 + \log m)$
- 2) $2\log 4 + \log m$
- 3) $\log 4 + 2\log m$
- 4) $\log 16 + 2\log m$


33 In which interval of $f(x) = \cos(x)$ is the inverse also a function?

- 1) $-\frac{\pi}{2} < x < \frac{\pi}{2}$
- 2) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$
- 3) $0 \leq x \leq \pi$
- 4) $\frac{\pi}{2} \leq x \leq \frac{3\pi}{2}$

34 When $x^2 + 3x - 4$ is subtracted from $x^3 + 3x^2 - 2x$, the difference is

- 1) $x^3 + 2x^2 - 5x + 4$
- 2) $x^3 + 2x^2 + x - 4$
- 3) $-x^3 + 4x^2 + x - 4$
- 4) $-x^3 - 2x^2 + 5x + 4$

35 Which equation is graphed in the diagram below?



- 1) $y = 3 \cos\left(\frac{\pi}{30}x\right) + 8$
- 2) $y = 3 \cos\left(\frac{\pi}{15}x\right) + 5$
- 3) $y = -3 \cos\left(\frac{\pi}{30}x\right) + 8$
- 4) $y = -3 \cos\left(\frac{\pi}{15}x\right) + 5$

36 The simplest form of $\frac{1 - \frac{4}{x}}{1 - \frac{2}{x} - \frac{8}{x^2}}$ is

- 1) $\frac{1}{2}$
- 2) $\frac{x}{x+2}$
- 3) $\frac{x}{3}$
- 4) $-\frac{x}{x-2}$

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

37 The solution set of the equation $\sqrt{2x-4} = x-2$ is

- 1) $\{-2, -4\}$
- 2) $\{2, 4\}$
- 3) $\{4\}$
- 4) $\{ \}$

38 There are eight people in a tennis club. Which expression can be used to find the number of different ways they can place first, second, and third in a tournament?

- 1) ${}_8P_3$
- 2) ${}_8C_3$
- 3) ${}_8P_5$
- 4) ${}_8C_5$


39 If $f(x) = 4x - x^2$ and $g(x) = \frac{1}{x}$, then $(f \circ g)\left(\frac{1}{2}\right)$ is

- equal to
- 1) $\frac{4}{7}$
 - 2) -2
 - 3) $\frac{7}{2}$
 - 4) 4

40 Which function is one-to-one?

- 1) $k(x) = x^2 + 2$
- 2) $g(x) = x^3 + 2$
- 3) $f(x) = |x| + 2$
- 4) $j(x) = x^4 + 2$

41 Which equation is represented by the graph below?



- 1) $(x-3)^2 + (y+1)^2 = 5$
- 2) $(x+3)^2 + (y-1)^2 = 5$
- 3) $(x-1)^2 + (y+3)^2 = 13$
- 4) $(x+3)^2 + (y-1)^2 = 13$





42 The conjugate of the complex expression $-5x + 4i$ is

- 1) $5x - 4i$
- 2) $5x + 4i$
- 3) $-5x - 4i$
- 4) $-5x + 4i$

43 A math club has 30 boys and 20 girls. Which expression represents the total number of different 5-member teams, consisting of 3 boys and 2 girls, that can be formed?


- 1) ${}_{30}P_3 \cdot {}_{20}P_2$
- 2) ${}_{30}C_3 \cdot {}_{20}C_2$
- 3) ${}_{30}P_3 + {}_{20}P_2$
- 4) ${}_{30}C_3 + {}_{20}C_2$

44 Which graph represents a function?



45 What is the graph of the solution set of

$$|2x - 1| > 5?$$



46 If $x = 3i$, $y = 2i$, and $z = m + i$, the expression xy^2z equals

- 1) $-12 - 12mi$
- 2) $-6 - 6mi$
- 3) $12 - 12mi$
- 4) $6 - 6mi$

47 Which calculator output shows the strongest linear relationship between x and y ?

Lin Reg

$$y = a + bx$$

$$a = 59.026$$

$$b = 6.767$$

- 1) $r = .8643$

Lin Reg

$$y = a + bx$$

$$a = .7$$

$$b = 24.2$$

- 2) $r = .8361$

Lin Reg

$$y = a + bx$$

$$a = 2.45$$

$$b = .95$$

- 3) $r = .6022$

Lin Reg

$$y = a + bx$$

$$a = -2.9$$

$$b = 24.1$$

- 4) $r = -.8924$

48 What is a positive value of $\tan \frac{1}{2}x$, when


$$\sin x = 0.8?$$

- 1) 0.5
- 2) 0.4
- 3) 0.33
- 4) 0.25

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

- 49 The expression $\sqrt[3]{64a^{16}}$ is equivalent to
- 1) $8a^4$
 - 2) $8a^8$
 - 3) $4a^5\sqrt[3]{a}$
 - 4) $4a\sqrt[3]{a^5}$
- 50 Which survey is *least* likely to contain bias?
- 1) surveying a sample of people leaving a movie theater to determine which flavor of ice cream is the most popular
 - 2) surveying the members of a football team to determine the most watched TV sport
 - 3) surveying a sample of people leaving a library to determine the average number of books a person reads in a year
 - 4) surveying a sample of people leaving a gym to determine the average number of hours a person exercises per week
- 51 The quantities p and q vary inversely. If $p = 20$ when $q = -2$, and $p = x$ when $q = -2x + 2$, then x equals
- 1) -4 and 5
 - 2) $\frac{20}{19}$
 - 3) -5 and 4
 - 4) $-\frac{1}{4}$
- 52 If $f(x) = \sqrt{9 - x^2}$, what are its domain and range?
- 1) domain: $\{x \mid -3 \leq x \leq 3\}$; range: $\{y \mid 0 \leq y \leq 3\}$
 - 2) domain: $\{x \mid x \neq \pm 3\}$; range: $\{y \mid 0 \leq y \leq 3\}$
 - 3) domain: $\{x \mid x \leq -3 \text{ or } x \geq 3\}$; range: $\{y \mid y \neq 0\}$
 - 4) domain: $\{x \mid x \neq 3\}$; range: $\{y \mid y \geq 0\}$
- 53 If the amount of time students work in any given week is normally distributed with a mean of 10 hours per week and a standard deviation of 2 hours, what is the probability a student works between 8 and 11 hours per week?
- 1) 34.1%
 - 2) 38.2%
 - 3) 53.2%
 - 4) 68.2%
- 54 What is the number of degrees in an angle whose measure is 2 radians?
- 1) $\frac{360}{\pi}$
 - 2) $\frac{\pi}{360}$
 - 3) 360
 - 4) 90
- 55 The graph below shows the average price of gasoline, in dollars, for the years 1997 to 2007.



What is the approximate range of this graph?

- 1) $1997 \leq x \leq 2007$
- 2) $1999 \leq x \leq 2007$
- 3) $0.97 \leq y \leq 2.38$
- 4) $1.27 \leq y \leq 2.38$

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

56 Approximately how many degrees does five radians equal?

- 1) 286
- 2) 900
- 3) $\frac{\pi}{36}$
- 4) 5π

57 Expressed with a rational denominator and in

simplest form, $\frac{x}{x - \sqrt{x}}$ is

- 1) $\frac{x^2 + x\sqrt{x}}{x^2 - x}$
- 2) $-\sqrt{x}$
- 3) $\frac{x + \sqrt{x}}{1 - x}$
- 4) $\frac{x + \sqrt{x}}{x - 1}$

58 The number of possible different 12-letter arrangements of the letters in the word "TRIGONOMETRY" is represented by

- 1) $\frac{12!}{3!}$
- 2) $\frac{12!}{6!}$
- 3) $\frac{{}^{12}P_{12}}{8}$
- 4) $\frac{{}^{12}P_{12}}{6!}$

59 In $\triangle PQR$, p equals

- 1) $\frac{r \sin P}{\sin Q}$
- 2) $\frac{r \sin P}{\sin R}$
- 3) $\frac{r \sin R}{\sin P}$
- 4) $\frac{q \sin R}{\sin Q}$

60 In $\triangle DEF$, $d = 5$, $e = 8$, and $m\angle D = 32$. How many distinct triangles can be drawn given these measurements?

- 1) 1
- 2) 2
- 3) 3
- 4) 0

61 If $\log_b x = 3 \log_b p - \left(2 \log_b t + \frac{1}{2} \log_b r \right)$, then the value of x is

- 1) $\frac{p^3}{\sqrt{t^2 r}}$
- 2) $p^3 t^2 r^{\frac{1}{2}}$
- 3) $\frac{p^3 t^2}{\sqrt{r}}$
- 4) $\frac{p^3}{t^2 \sqrt{r}}$

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

62 The value of $\sin(180 + x)$ is equivalent to

- 1) $-\sin x$
- 2) $-\sin(90 - x)$
- 3) $\sin x$
- 4) $\sin(90 - x)$

63 The area of triangle ABC is 42. If $AB = 8$ and $m\angle B = 61$, the length of BC is approximately

- 1) 5.1
- 2) 9.2
- 3) 12.0
- 4) 21.7

64 The expression $(2 - 3\sqrt{x})^2$ is equivalent to

- 1) $4 - 9x$
- 2) $4 - 3x$
- 3) $4 - 12\sqrt{x} + 9x$
- 4) $4 - 12\sqrt{x} + 6x$

65 A spinner is divided into eight equal sections. Five sections are red and three are green. If the spinner is spun three times, what is the probability that it lands on red *exactly* twice?

- 1) $\frac{25}{64}$
- 2) $\frac{45}{512}$
- 3) $\frac{75}{512}$
- 4) $\frac{225}{512}$


66 Which expression always equals 1?

- 1) $\cos^2 x - \sin^2 x$
- 2) $\cos^2 x + \sin^2 x$
- 3) $\cos x - \sin x$
- 4) $\cos x + \sin x$

67 What is the common difference in the sequence $2a + 1, 4a + 4, 6a + 7, 8a + 10, \dots$?

- 1) $2a + 3$
- 2) $-2a - 3$
- 3) $2a + 5$
- 4) $-2a + 5$

68 Which diagram represents a relation that is both one-to-one and onto?



Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

- 69 What is the fourth term in the binomial expansion $(x - 2)^8$?
- 1) $448x^5$
 - 2) $448x^4$
 - 3) $-448x^5$
 - 4) $-448x^4$
- 70 What is the common ratio of the geometric sequence shown below?
 $-2, 4, -8, 16, \dots$
- 1) $-\frac{1}{2}$
 - 2) 2
 - 3) -2
 - 4) -6
- 71 In the interval $0^\circ \leq x < 360^\circ$, $\tan x$ is undefined when x equals
- 1) 0° and 90°
 - 2) 90° and 180°
 - 3) 180° and 270°
 - 4) 90° and 270°
- 72 Expressed in simplest form, $\sqrt{-18} - \sqrt{-32}$ is
- 1) $-\sqrt{2}$
 - 2) $-7\sqrt{2}$
 - 3) $-i\sqrt{2}$
 - 4) $7i\sqrt{2}$
- 73 Which statement about the graph of the equation $y = e^x$ is *not* true?
- 1) It is asymptotic to the x -axis.
 - 2) The domain is the set of all real numbers.
 - 3) It lies in Quadrants I and II.
 - 4) It passes through the point $(e, 1)$.
- 74 Which step can be used when solving $x^2 - 6x - 25 = 0$ by completing the square?
- 1) $x^2 - 6x + 9 = 25 + 9$
 - 2) $x^2 - 6x - 9 = 25 - 9$
 - 3) $x^2 - 6x + 36 = 25 + 36$
 - 4) $x^2 - 6x - 36 = 25 - 36$
- 75 In $\triangle MNP$, $m = 6$ and $n = 10$. Two distinct triangles can be constructed if the measure of angle M is
- 1) 35
 - 2) 40
 - 3) 45
 - 4) 50
- 76 Which ordered pair is a solution of the system of equations shown below? $x + y = 5$
 $(x + 3)^2 + (y - 3)^2 = 53$
- 1) (2, 3)
 - 2) (5, 0)
 - 3) (-5, 10)
 - 4) (-4, 9)

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org


77 Which value of r represents data with a strong positive linear correlation between two variables?

- 1) 0.89
- 2) 0.34
- 3) 1.04
- 4) 0.01

78 What is the product of the roots of the quadratic equation $2x^2 - 7x = 5$?

- 1) 5
- 2) $\frac{5}{2}$
- 3) -5
- 4) $-\frac{5}{2}$

79 In the right triangle shown below, what is the measure of angle S , to the *nearest minute*?



- 1) $28^\circ 1'$
- 2) $28^\circ 4'$
- 3) $61^\circ 56'$
- 4) $61^\circ 93'$

80 What are the coordinates of the center of a circle whose equation is $x^2 + y^2 - 16x + 6y + 53 = 0$?

- 1) $(-8, -3)$
- 2) $(-8, 3)$
- 3) $(8, -3)$
- 4) $(8, 3)$

81 Which equation has roots with the sum equal to $\frac{9}{4}$ and the product equal to $\frac{3}{4}$?

- 1) $4x^2 + 9x + 3 = 0$
- 2) $4x^2 + 9x - 3 = 0$
- 3) $4x^2 - 9x + 3 = 0$
- 4) $4x^2 - 9x - 3 = 0$

82 The expression $4 + \sum_{k=2}^5 3(k-x)$ is equal to

- 1) $58 - 4x$
- 2) $46 - 4x$
- 3) $58 - 12x$
- 4) $46 - 12x$

83 The table below displays the results of a survey regarding the number of pets each student in a class has. The average number of pets per student in this class is 2.

Number of Pets	0	1	2	3	4	5
Number of Students	4	6	10	0	k	2

What is the value of k for this table?

- 1) 9
- 2) 2
- 3) 8
- 4) 4

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

- 84 What are the sum and product of the roots of the equation $6x^2 - 4x - 12 = 0$?
- 1) sum = $-\frac{2}{3}$; product = -2
 - 2) sum = $\frac{2}{3}$; product = -2
 - 3) sum = -2 ; product = $\frac{2}{3}$
 - 4) sum = -2 ; product = $-\frac{2}{3}$
- 85 When factored completely, the expression $3x^3 - 5x^2 - 48x + 80$ is equivalent to
- 1) $(x^2 - 16)(3x - 5)$
 - 2) $(x^2 + 16)(3x - 5)(3x + 5)$
 - 3) $(x + 4)(x - 4)(3x - 5)$
 - 4) $(x + 4)(x - 4)(3x - 5)(3x - 5)$
- 86 If $2x^3 = y$, then $\log y$ equals
- 1) $\log(2x) + \log 3$
 - 2) $3 \log(2x)$
 - 3) $3 \log 2 + 3 \log x$
 - 4) $\log 2 + 3 \log x$
- 87 The expression $\frac{1}{7 - \sqrt{11}}$ is equivalent to
- 1) $\frac{7 + \sqrt{11}}{38}$
 - 2) $\frac{7 - \sqrt{11}}{38}$
 - 3) $\frac{7 + \sqrt{11}}{60}$
 - 4) $\frac{7 - \sqrt{11}}{60}$
- 88 How many different six-letter arrangements can be made using the letters of the word "TATTOO"?
- 1) 60
 - 2) 90
 - 3) 120
 - 4) 720
- 89 For $y = \frac{3}{\sqrt{x-4}}$, what are the domain and range?
- 1) $\{x|x > 4\}$ and $\{y|y > 0\}$
 - 2) $\{x|x \geq 4\}$ and $\{y|y > 0\}$
 - 3) $\{x|x > 4\}$ and $\{y|y \geq 0\}$
 - 4) $\{x|x \geq 4\}$ and $\{y|y \geq 0\}$
- 90 If $\sin \theta < 0$ and $\cot \theta > 0$, in which quadrant does the terminal side of angle θ lie?
- 1) I
 - 2) II
 - 3) III
 - 4) IV
- 91 Which expression is equivalent to $(9x^2y^6)^{-\frac{1}{2}}$?
- 1) $\frac{1}{3xy^3}$
 - 2) $3xy^3$
 - 3) $\frac{3}{xy^3}$
 - 4) $\frac{xy^3}{3}$

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

- 92 Given $\triangle ABC$ with $a = 9$, $b = 10$, and $m\angle B = 70$, what type of triangle can be drawn?
- 1) an acute triangle, only
 - 2) an obtuse triangle, only
 - 3) both an acute triangle and an obtuse triangle
 - 4) neither an acute triangle nor an obtuse triangle

- 93 The value of $\csc 138^\circ 23'$ rounded to four decimal places is
- 1) -1.3376
 - 2) -1.3408
 - 3) 1.5012
 - 4) 1.5057

- 94 Which function is one-to-one?
- 1) $f(x) = |x|$
 - 2) $f(x) = 2^x$
 - 3) $f(x) = x^2$
 - 4) $f(x) = \sin x$

- 95 What is the solution set of the equation $\frac{30}{x^2 - 9} + 1 = \frac{5}{x - 3}$?
- 1) $\{2, 3\}$
 - 2) $\{2\}$
 - 3) $\{3\}$
 - 4) $\{ \}$





- 96 If $f(x) = 2x^2 - 3x + 1$ and $g(x) = x + 5$, what is $f(g(x))$?
- 1) $2x^2 + 17x + 36$
 - 2) $2x^2 + 17x + 66$
 - 3) $2x^2 - 3x + 6$
 - 4) $2x^2 - 3x + 36$

- 97 The sum of $\sqrt[3]{6a^4b^2}$ and $\sqrt[3]{162a^4b^2}$, expressed in simplest radical form, is
- 1) $\sqrt[6]{168a^8b^4}$
 - 2) $2a^2b^3\sqrt[3]{21a^2b}$
 - 3) $4a^3\sqrt[3]{6ab^2}$
 - 4) $10a^2b^3\sqrt[3]{8}$

- 98 The expression $\frac{x^2 + 9x - 22}{x^2 - 121} \div (2 - x)$ is equivalent to
- 1) $x - 11$
 - 2) $\frac{1}{x - 11}$
 - 3) $11 - x$
 - 4) $\frac{1}{11 - x}$

- 99 Which graph represents the solution set of

$$\left| \frac{4x - 5}{3} \right| > 1?$$

- 1) 
- 2) 
- 3) 
- 4) 

100 Which statement regarding the inverse function is true?

- 1) A domain of $y = \sin^{-1}x$ is $[0, 2\pi]$.
- 2) The range of $y = \sin^{-1}x$ is $[-1, 1]$.
- 3) A domain of $y = \cos^{-1}x$ is $(-\infty, \infty)$.
- 4) The range of $y = \cos^{-1}x$ is $[0, \pi]$.

101 Which summation represents

$5 + 7 + 9 + 11 + \dots + 43$?

- 1) $\sum_{n=5}^{43} n$
- 2) $\sum_{n=1}^{20} (2n + 3)$
- 3) $\sum_{n=4}^{24} (2n - 3)$
- 4) $\sum_{n=3}^{23} (3n - 4)$

102 The sum of the first eight terms of the series


$3 - 12 + 48 - 192 + \dots$ is

- 1) $-13, 107$
- 2) $-21, 845$
- 3) $-39, 321$
- 4) $-65, 535$

103 The expression $x^2(x + 2) - (x + 2)$ is equivalent to

- 1) x^2
- 2) $x^2 - 1$
- 3) $x^3 + 2x^2 - x + 2$
- 4) $(x + 1)(x - 1)(x + 2)$

104 What is the range of the function shown below?



- 1) $x \leq 0$
- 2) $x \geq 0$
- 3) $y \leq 0$
- 4) $y \geq 0$

105 Which expression is equivalent to $\frac{x^{-1}y^4}{3x^{-5}y^{-1}}$?

- 1) $\frac{x^4y^5}{3}$
- 2) $\frac{x^5y^4}{3}$
- 3) $3x^4y^5$
- 4) $\frac{y^4}{3x^5}$

106 The points $(2, 3)$, $(4, \frac{3}{4})$, and $(6, d)$ lie on the graph of a function. If y is inversely proportional to the square of x , what is the value of d ?

- 1) 1
- 2) $\frac{1}{3}$
- 3) 3
- 4) 27

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

- 107 What is the common difference of the arithmetic sequence below?

$$-7x, -4x, -x, 2x, 5x, \dots$$

- 1) -3
- 2) $-3x$
- 3) 3
- 4) $3x$

- 108 If p varies inversely as q , and $p = 10$ when $q = \frac{3}{2}$,

what is the value of p when $q = \frac{3}{5}$?

- 1) 25
- 2) 15
- 3) 9
- 4) 4

- 109 If $\sin A = -\frac{7}{25}$ and $\angle A$ terminates in Quadrant IV,

$\tan A$ equals

- 1) $-\frac{7}{25}$
- 2) $-\frac{7}{24}$
- 3) $-\frac{24}{7}$
- 4) $-\frac{24}{25}$

- 110 A population of rabbits doubles every 60 days

according to the formula $P = 10(2)^{\frac{t}{60}}$, where P is the population of rabbits on day t . What is the value of t when the population is 320?

- 1) 240
- 2) 300
- 3) 660
- 4) 960

- 111 What is the sum of the first 19 terms of the sequence 3, 10, 17, 24, 31, ...?

- 1) 1188
- 2) 1197
- 3) 1254
- 4) 1292

- 112 In a certain high school, a survey revealed the mean amount of bottled water consumed by students each day was 153 bottles with a standard deviation of 22 bottles. Assuming the survey represented a normal distribution, what is the range of the number of bottled waters that approximately 68.2% of the students drink?

- 1) 131 – 164
- 2) 131 – 175
- 3) 142 – 164
- 4) 142 – 175


- 113 Which expression is equivalent to $\sum_{n=1}^4 (a-n)^2$?

- 1) $2a^2 + 17$
- 2) $4a^2 + 30$
- 3) $2a^2 - 10a + 17$
- 4) $4a^2 - 20a + 30$

- 114 Two sides of a triangular-shaped sandbox measure 22 feet and 13 feet. If the angle between these two sides measures 55° , what is the area of the sandbox, to the nearest square foot?

- 1) 82
- 2) 117
- 3) 143
- 4) 234

115 Which equation is represented by the graph below?



- 1) $y = 2 \cos 3x$
- 2) $y = 2 \sin 3x$
- 3) $y = 2 \cos \frac{2\pi}{3} x$
- 4) $y = 2 \sin \frac{2\pi}{3} x$

116 The formula to determine continuously compounded interest is $A = Pe^{rt}$, where A is the amount of money in the account, P is the initial investment, r is the interest rate, and t is the time, in years. Which equation could be used to determine the value of an account with an \$18,000 initial investment, at an interest rate of 1.25% for 24 months?

- 1) $A = 18,000e^{1.25 \cdot 2}$
- 2) $A = 18,000e^{1.25 \cdot 24}$
- 3) $A = 18,000e^{0.0125 \cdot 2}$
- 4) $A = 18,000e^{0.0125 \cdot 24}$

117 What is the product of the roots of $x^2 - 4x + k = 0$ if one of the roots is 7?

- 1) 21
- 2) -11
- 3) -21
- 4) -77

118 Which expression is equivalent to $\frac{2x^{-2}y^{-2}}{4y^{-5}}$?

- 1) $\frac{y^3}{2x^2}$
- 2) $\frac{2y^3}{x^2}$
- 3) $\frac{2x^2}{y^3}$
- 4) $\frac{x^2}{2y^3}$

119 What is the product of $\left(\frac{x}{4} - \frac{1}{3}\right)$ and $\left(\frac{x}{4} + \frac{1}{3}\right)$?

- 1) $\frac{x^2}{8} - \frac{1}{9}$
- 2) $\frac{x^2}{16} - \frac{1}{9}$
- 3) $\frac{x^2}{8} - \frac{x}{6} - \frac{1}{9}$
- 4) $\frac{x^2}{16} - \frac{x}{6} - \frac{1}{9}$


120 Which problem involves evaluating ${}_6P_4$?

- 1) How many different four-digit ID numbers can be formed using 1, 2, 3, 4, 5, and 6 without repetition?
- 2) How many different subcommittees of four can be chosen from a committee having six members?
- 3) How many different outfits can be made using six shirts and four pairs of pants?
- 4) How many different ways can one boy and one girl be selected from a group of four boys and six girls?

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

- 121 In the diagram below, the length of which line segment is equal to the exact value of $\sin \theta$?



- 1) \overline{TO}
 - 2) \overline{TS}
 - 3) \overline{OR}
 - 4) \overline{OS}
- 122 Theresa is comparing the graphs of $y = 2^x$ and $y = 5^x$. Which statement is true?
- 1) The y-intercept of $y = 2^x$ is $(0, 2)$, and the y-intercept of $y = 5^x$ is $(0, 5)$.
 - 2) Both graphs have a y-intercept of $(0, 1)$, and $y = 2^x$ is steeper for $x > 0$.
 - 3) Both graphs have a y-intercept of $(0, 1)$, and $y = 5^x$ is steeper for $x > 0$.
 - 4) Neither graph has a y-intercept.

- 123 What is the product of $\left(\frac{2}{5}x - \frac{3}{4}y^2\right)$ and

$$\left(\frac{2}{5}x + \frac{3}{4}y^2\right)?$$

- 1) $\frac{4}{25}x^2 - \frac{9}{16}y^4$
 - 2) $\frac{4}{25}x - \frac{9}{16}y^2$
 - 3) $\frac{2}{5}x^2 - \frac{3}{4}y^4$
 - 4) $\frac{4}{5}x$
- 124 What is the solution set for $2\cos \theta - 1 = 0$ in the interval $0^\circ \leq \theta < 360^\circ$?
- 1) $\{30^\circ, 150^\circ\}$
 - 2) $\{60^\circ, 120^\circ\}$
 - 3) $\{30^\circ, 330^\circ\}$
 - 4) $\{60^\circ, 300^\circ\}$

- 125 Given the relation $\{(8, 2), (3, 6), (7, 5), (k, 4)\}$, which value of k will result in the relation *not* being a function?
- 1) 1
 - 2) 2
 - 3) 3
 - 4) 4

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

126 A study finds that 80% of the local high school students text while doing homework. Ten students are selected at random from the local high school. Which expression would be part of the process used to determine the probability that, *at most*, 7 of the 10 students text while doing homework?

- 1) ${}_{10}C_6 \left(\frac{4}{5}\right)^6 \left(\frac{1}{5}\right)^4$
- 2) ${}_{10}C_7 \left(\frac{4}{5}\right)^{10} \left(\frac{1}{5}\right)^7$
- 3) ${}_{10}C_8 \left(\frac{7}{10}\right)^{10} \left(\frac{3}{10}\right)^2$
- 4) ${}_{10}C_9 \left(\frac{7}{10}\right)^9 \left(\frac{3}{10}\right)^1$

127 If $\log 2 = a$ and $\log 3 = b$, the expression $\log \frac{9}{20}$ is equivalent to

- 1) $2b - a + 1$
- 2) $2b - a - 1$
- 3) $b^2 - a + 10$
- 4) $\frac{2b}{a + 1}$

128 Which expression represents the third term in the expansion of $(2x^4 - y)^3$?

- 1) $-y^3$
- 2) $-6x^4y^2$
- 3) $6x^4y^2$
- 4) $2x^4y^2$

129 The expression $(x + i)^2 - (x - i)^2$ is equivalent to

- 1) 0
- 2) -2
- 3) $-2 + 4xi$
- 4) $4xi$

130 As shown in the table below, a person's target heart rate during exercise changes as the person gets older.

Age (years)	Target Heart Rate (beats per minute)
20	135
25	132
30	129
35	125
40	122
45	119
50	115

Which value represents the linear correlation coefficient, rounded to the *nearest thousandth*, between a person's age, in years, and that person's target heart rate, in beats per minute?

- 1) -0.999
- 2) -0.664
- 3) 0.998
- 4) 1.503

- 131 What is the period of the graph $y = \frac{1}{2} \sin 6x$?
- 1) $\frac{\pi}{6}$
 - 2) $\frac{\pi}{3}$
 - 3) $\frac{\pi}{2}$
 - 4) 6π
- 132 If $\sin x = \sin y = a$ and $\cos x = \cos y = b$, then $\cos(x - y)$ is
- 1) $b^2 - a^2$
 - 2) $b^2 + a^2$
 - 3) $2b - 2a$
 - 4) $2b + 2a$
- 133 When $x^{-1} + 1$ is divided by $x + 1$, the quotient equals
- 1) 1
 - 2) $\frac{1}{x}$
 - 3) x
 - 4) $-\frac{1}{x}$
- 134 If $\tan\left(\arccos \frac{\sqrt{3}}{k}\right) = \frac{\sqrt{3}}{3}$, then k is
- 1) 1
 - 2) 2
 - 3) $\sqrt{2}$
 - 4) $3\sqrt{2}$
- 135 Which expression is equivalent to $(3x^2)^{-1}$?
- 1) $\frac{1}{3x^2}$
 - 2) $-3x^2$
 - 3) $\frac{1}{9x^2}$
 - 4) $-9x^2$
- 136 The yearbook staff has designed a survey to learn student opinions on how the yearbook could be improved for this year. If they want to distribute this survey to 100 students and obtain the most reliable data, they should survey
- 1) every third student sent to the office
 - 2) every third student to enter the library
 - 3) every third student to enter the gym for the basketball game
 - 4) every third student arriving at school in the morning
- 137 What is the solution set of the equation $3x^5 - 48x = 0$?
- 1) $\{0, \pm 2\}$
 - 2) $\{0, \pm 2, 3\}$
 - 3) $\{0, \pm 2, \pm 2i\}$
 - 4) $\{\pm 2, \pm 2i\}$
- 138 If order does *not* matter, which selection of students would produce the most possible committees?
- 1) 5 out of 15
 - 2) 5 out of 25
 - 3) 20 out of 25
 - 4) 15 out of 25

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions





www.jmap.org

139 Which value of k satisfies the equation

$$8^{3k+4} = 4^{2k-1}?$$

- 1) -1
- 2) $-\frac{9}{4}$
- 3) -2
- 4) $-\frac{14}{5}$

140 Which sketch shows the inverse of $y = a^x$, where $a > 1$?

- 1) 
- 2) 
- 3) 
- 4) 

141 If \$5000 is invested at a rate of 3% interest compounded quarterly, what is the value of the investment in 5 years? (Use the formula

$$A = P \left(1 + \frac{r}{n} \right)^{nt},$$

where A is the amount accrued, P

is the principal, r is the interest rate, n is the number of times per year the money is compounded, and t is the length of time, in years.)





- 1) \$5190.33
- 2) \$5796.37
- 3) \$5805.92
- 4) \$5808.08

142 Which expression is equivalent to $(n \circ m \circ p)(x)$, given $m(x) = \sin x$, $n(x) = 3x$, and $p(x) = x^2$?

- 1) $\sin(3x)^2$
- 2) $3 \sin x^2$
- 3) $\sin^2(3x)$
- 4) $3 \sin^2 x$

143 Which graph represents the solution set of

$$\frac{x+16}{x-2} \leq 7?$$

- 1) 
- 2) 
- 3) 
- 4) 

144 The expression $\frac{\cot x}{\csc x}$ is equivalent to

- 1) $\sin x$
- 2) $\cos x$
- 3) $\tan x$
- 4) $\sec x$

145 The relationship between t , a student's test scores, and d , the student's success in college, is modeled by the equation $d = 0.48t + 75.2$. Based on this linear regression model, the correlation coefficient could be

- 1) between -1 and 0
- 2) between 0 and 1
- 3) equal to -1
- 4) equal to 0

146 Which ordered pair is in the solution set of the system of equations shown below?

$$y^2 - x^2 + 32 = 0$$

$$3y - x = 0$$

- 1) $(2, 6)$
- 2) $(3, 1)$
- 3) $(-1, -3)$
- 4) $(-6, -2)$


147 An angle, P , drawn in standard position, terminates in Quadrant II if

- 1) $\cos P < 0$ and $\csc P < 0$
- 2) $\sin P > 0$ and $\cos P > 0$
- 3) $\csc P > 0$ and $\cot P < 0$
- 4) $\tan P < 0$ and $\sec P > 0$

148 The expression $\left(\sqrt[3]{27x^2}\right)\left(\sqrt[3]{16x^4}\right)$ is equivalent

- to
- 1) $12x^2\sqrt[3]{2}$
 - 2) $12x^3\sqrt[3]{2x}$
 - 3) $6x^3\sqrt[3]{2x^2}$
 - 4) $6x^2\sqrt[3]{2}$

149 What is the domain of the function shown below?



- 1) $-1 \leq x \leq 6$
- 2) $-1 \leq y \leq 6$
- 3) $-2 \leq x \leq 5$
- 4) $-2 \leq y \leq 5$

150 Which relation is both one-to-one and onto?

- 1)
- 2)
- 3)
- 4)

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

151 If $\sin A = \frac{1}{3}$, what is the value of $\cos 2A$?

- 1) $-\frac{2}{3}$
- 2) $\frac{2}{3}$
- 3) $-\frac{7}{9}$
- 4) $\frac{7}{9}$

152 Which expression, when rounded to three decimal places, is equal to -1.155 ?

- 1) $\sec\left(\frac{5\pi}{6}\right)$
- 2) $\tan(49^\circ 20')$
- 3) $\sin\left(-\frac{3\pi}{5}\right)$
- 4) $\csc(-118^\circ)$

153 If d varies inversely as t , and $d = 20$ when $t = 2$, what is the value of t when $d = -5$?

- 1) 8
- 2) 2
- 3) -8
- 4) -2

154 Which equation represents a graph that has a period of 4π ?

- 1) $y = 3 \sin \frac{1}{2}x$
- 2) $y = 3 \sin 2x$
- 3) $y = 3 \sin \frac{1}{4}x$
- 4) $y = 3 \sin 4x$

155 What is the common ratio of the sequence

$$\frac{1}{64}a^5b^3, -\frac{3}{32}a^3b^4, \frac{9}{16}ab^5, \dots?$$

- 1) $-\frac{3b}{2a^2}$
- 2) $-\frac{6b}{a^2}$
- 3) $-\frac{3a^2}{b}$
- 4) $-\frac{6a^2}{b}$

156 Given angle A in Quadrant I with $\sin A = \frac{12}{13}$ and

angle B in Quadrant II with $\cos B = -\frac{3}{5}$, what is the value of $\cos(A - B)$?

- 1) $\frac{33}{65}$
- 2) $-\frac{33}{65}$
- 3) $\frac{63}{65}$
- 4) $-\frac{63}{65}$

157 What is the conjugate of $\frac{1}{2} + \frac{3}{2}i$?

- 1) $-\frac{1}{2} + \frac{3}{2}i$
- 2) $\frac{1}{2} - \frac{3}{2}i$
- 3) $\frac{3}{2} + \frac{1}{2}i$
- 4) $-\frac{1}{2} - \frac{3}{2}i$

158 What is the middle term in the expansion of

$$\left(\frac{x}{2} - 2y\right)^6?$$

- 1) $20x^3y^3$
- 2) $-\frac{15}{4}x^4y^2$
- 3) $-20x^3y^3$
- 4) $\frac{15}{4}x^4y^2$

159 Susie invests \$500 in an account that is compounded continuously at an annual interest rate of 5%, according to the formula $A = Pe^{rt}$, where A is the amount accrued, P is the principal, r is the rate of interest, and t is the time, in years.

Approximately how many years will it take for Susie's money to double?

- 1) 1.4
- 2) 6.0
- 3) 13.9
- 4) 14.7

160 In $\triangle KLM$, $KL = 20$, $LM = 13$, and $m\angle K = 40$. The measure of $\angle M$?


- 1) must be between 0° and 90°
- 2) must equal 90°
- 3) must be between 90° and 180°
- 4) is ambiguous

161 How many negative solutions to the equation

$$2x^3 - 4x^2 + 3x - 1 = 0$$
 exist?

- 1) 1
- 2) 2
- 3) 3
- 4) 0

162 Which graph represents the function $\log_2 x = y$?



Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

163 If $x^2 + 2 = 6x$ is solved by completing the square, an intermediate step would be

- 1) $(x + 3)^2 = 7$
- 2) $(x - 3)^2 = 7$
- 3) $(x - 3)^2 = 11$
- 4) $(x - 6)^2 = 34$

164 The solutions of the equation $y^2 - 3y = 9$ are

- 1) $\frac{3 \pm 3i\sqrt{3}}{2}$
- 2) $\frac{3 \pm 3i\sqrt{5}}{2}$
- 3) $\frac{-3 \pm 3\sqrt{5}}{2}$
- 4) $\frac{3 \pm 3\sqrt{5}}{2}$

165 Which relation is *not* a function?

- 1) $(x - 2)^2 + y^2 = 4$
- 2) $x^2 + 4x + y = 4$
- 3) $x + y = 4$
- 4) $xy = 4$

166 The product of i^7 and i^5 is equivalent to

- 1) 1
- 2) -1
- 3) i
- 4) $-i$

167 In which graph is θ coterminal with an angle of -70° ?

1)

2)

3)

4)

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

168 If $f(x) = \frac{x}{x^2 - 16}$, what is the value of $f(-10)$?

- 1) $-\frac{5}{2}$
- 2) $-\frac{5}{42}$
- 3) $\frac{5}{58}$
- 4) $\frac{5}{18}$

169 A circle has a radius of 4 inches. In inches, what is the length of the arc intercepted by a central angle of 2 radians?

- 1) 2π
- 2) 2
- 3) 8π
- 4) 8

170 In $\triangle ABC$, $m\angle A = 74$, $a = 59.2$, and $c = 60.3$. What are the two possible values for $m\angle C$, to the nearest tenth?

- 1) 73.7 and 106.3
- 2) 73.7 and 163.7
- 3) 78.3 and 101.7
- 4) 78.3 and 168.3

171 The equation $y - 2\sin\theta = 3$ may be rewritten as

- 1) $f(y) = 2\sin x + 3$
- 2) $f(y) = 2\sin\theta + 3$
- 3) $f(x) = 2\sin\theta + 3$
- 4) $f(\theta) = 2\sin\theta + 3$


172 The solution set of the inequality $x^2 - 3x > 10$ is

- 1) $\{x | -2 < x < 5\}$
- 2) $\{x | 0 < x < 3\}$
- 3) $\{x | x < -2 \text{ or } x > 5\}$
- 4) $\{x | x < -5 \text{ or } x > 2\}$

173 The value of x in the equation $4^{2x+5} = 8^{3x}$ is

- 1) 1
- 2) 2
- 3) 5
- 4) -10

174 Which graph represents the solution set of $|6x - 7| \leq 5$?



175 In simplest form, $\sqrt{-300}$ is equivalent to

- 1) $3i\sqrt{10}$
- 2) $5i\sqrt{12}$
- 3) $10i\sqrt{3}$
- 4) $12i\sqrt{5}$

176 What is the fifteenth term of the geometric sequence $-\sqrt{5}, \sqrt{10}, -2\sqrt{5}, \dots$?

- 1) $-128\sqrt{5}$
- 2) $128\sqrt{10}$
- 3) $-16384\sqrt{5}$
- 4) $16384\sqrt{10}$

177 What are the values of θ in the interval $0^\circ \leq \theta < 360^\circ$ that satisfy the equation

$$\tan \theta - \sqrt{3} = 0?$$

- 1) $60^\circ, 240^\circ$
- 2) $72^\circ, 252^\circ$
- 3) $72^\circ, 108^\circ, 252^\circ, 288^\circ$
- 4) $60^\circ, 120^\circ, 240^\circ, 300^\circ$





178 The solution set of $\sqrt{3x+16} = x+2$ is

- 1) $\{-3, 4\}$
- 2) $\{-4, 3\}$
- 3) $\{3\}$
- 4) $\{-4\}$

179 The minimum point on the graph of the equation $y = f(x)$ is $(-1, -3)$. What is the minimum point on the graph of the equation $y = f(x) + 5$?

- 1) $(-1, 2)$
- 2) $(-1, -8)$
- 3) $(4, -3)$
- 4) $(-6, -3)$

180 On January 1, a share of a certain stock cost \$180. Each month thereafter, the cost of a share of this stock decreased by one-third. If x represents the time, in months, and y represents the cost of the stock, in dollars, which graph best represents the cost of a share over the following 5 months?



181 The expression $2i^2 + 3i^3$ is equivalent to

- 1) $-2 - 3i$
- 2) $2 - 3i$
- 3) $-2 + 3i$
- 4) $2 + 3i$

182 The expression $\frac{a^2b^{-3}}{a^{-4}b^2}$ is equivalent to

- 1) $\frac{a^6}{b^5}$
- 2) $\frac{b^5}{a^6}$
- 3) $\frac{a^2}{b}$
- 4) $a^{-2}b^{-1}$

183 A four-digit serial number is to be created from the digits 0 through 9. How many of these serial numbers can be created if 0 can *not* be the first digit, no digit may be repeated, and the last digit must be 5?


- 1) 448
- 2) 504
- 3) 2,240
- 4) 2,520

184 When simplified, the expression $\left(\frac{w^{-5}}{w^{-9}}\right)^{\frac{1}{2}}$ is

equivalent to

- 1) w^{-7}
- 2) w^2
- 3) w^7
- 4) w^{14}

185 The graph of $y = x^3 - 4x^2 + x + 6$ is shown below.



What is the product of the roots of the equation

$$x^3 - 4x^2 + x + 6 = 0?$$

- 1) -36
- 2) -6
- 3) 6
- 4) 4

186 What is the radian measure of the smaller angle formed by the hands of a clock at 7 o'clock?

- 1) $\frac{\pi}{2}$
- 2) $\frac{2\pi}{3}$
- 3) $\frac{5\pi}{6}$
- 4) $\frac{7\pi}{6}$

187 The expression $(3 - 7i)^2$ is equivalent to

- 1) $-40 + 0i$
- 2) $-40 - 42i$
- 3) $58 + 0i$
- 4) $58 - 42i$


188 Which values of x are in the solution set of the following system of equations?

$$y = 3x - 6$$

$$y = x^2 - x - 6$$

- 1) $0, -4$
- 2) $0, 4$
- 3) $6, -2$
- 4) $-6, 2$

189 In the diagram below of right triangle JTM , $JT = 12$, $JM = 6$, and $m\angle JMT = 90$.




What is the value of $\cot J$?

- 1) $\frac{\sqrt{3}}{3}$
- 2) 2
- 3) $\sqrt{3}$
- 4) $\frac{2\sqrt{3}}{3}$

190 What is the range of $f(x) = (x + 4)^2 + 7$?

- 1) $y \geq -4$
- 2) $y \geq 4$
- 3) $y = 7$
- 4) $y \geq 7$

191 Which equation is sketched in the diagram below?




- 1) $y = \csc x$
- 2) $y = \sec x$
- 3) $y = \cot x$
- 4) $y = \tan x$

192 What is the conjugate of $-2 + 3i$?

- 1) $-3 + 2i$
- 2) $-2 - 3i$
- 3) $2 - 3i$
- 4) $3 + 2i$

193 The graph of $y = f(x)$ is shown below.



Which set lists all the real solutions of $f(x) = 0$?

- 1) $\{-3, 2\}$
- 2) $\{-2, 3\}$
- 3) $\{-3, 0, 2\}$
- 4) $\{-2, 0, 3\}$

194 What is the value of x in the equation





$$9^{3x+1} = 27^{x+2}?$$

- 1) 1
- 2) $\frac{1}{3}$
- 3) $\frac{1}{2}$
- 4) $\frac{4}{3}$


195 The solution set of $4^{x^2+4x} = 2^{-6}$ is

- 1) $\{1, 3\}$
- 2) $\{-1, 3\}$
- 3) $\{-1, -3\}$
- 4) $\{1, -3\}$





196 Which graph represents a relation that is *not* a function?




197 The graph below shows the function $f(x)$.



Which graph represents the function $f(x + 2)$?



198 Which ratio represents $\csc A$ in the diagram below?



- 1) $\frac{25}{24}$
- 2) $\frac{25}{7}$
- 3) $\frac{24}{7}$
- 4) $\frac{7}{24}$

199 The lengths of 100 pipes have a normal distribution with a mean of 102.4 inches and a standard deviation of 0.2 inch. If one of the pipes measures exactly 102.1 inches, its length lies

- 1) below the 16th percentile
- 2) between the 50th and 84th percentiles
- 3) between the 16th and 50th percentiles
- 4) above the 84th percentile

200 What is the period of the function

$$y = \frac{1}{2} \sin\left(\frac{x}{3} - \pi\right)?$$

- 1) $\frac{1}{2}$
- 2) $\frac{1}{3}$
- 3) $\frac{2}{3}\pi$
- 4) 6π

201 The expression $x^{-\frac{2}{5}}$ is equivalent to

- 1) $-\sqrt[2]{x^5}$
- 2) $-\sqrt[5]{x^2}$
- 3) $\frac{1}{\sqrt[2]{x^5}}$
- 4) $\frac{1}{\sqrt[5]{x^2}}$


202 The roots of the equation $2x^2 + 7x - 3 = 0$ are

- 1) $-\frac{1}{2}$ and -3
- 2) $\frac{1}{2}$ and 3
- 3) $\frac{-7 \pm \sqrt{73}}{4}$
- 4) $\frac{7 \pm \sqrt{73}}{4}$


203 If $\sin A = \frac{2}{3}$ where $0^\circ < A < 90^\circ$, what is the value of $\sin 2A$?

- 1) $\frac{2\sqrt{5}}{3}$
- 2) $\frac{2\sqrt{5}}{9}$
- 3) $\frac{4\sqrt{5}}{9}$
- 4) $-\frac{4\sqrt{5}}{9}$


204 Which graph shows $y = \cos^{-1}x$?




1)



2)



3)



4)


Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org


205 The conjugate of $7 - 5i$ is

- 1) $-7 - 5i$
- 2) $-7 + 5i$
- 3) $7 - 5i$
- 4) $7 + 5i$


206 Which graph represents one complete cycle of the equation $y = \sin 3\pi x$?




1)



2)



3)



4)

207 Twenty different cameras will be assigned to several boxes. Three cameras will be randomly selected and assigned to box A. Which expression can be used to calculate the number of ways that three cameras can be assigned to box A?

- 1) $20!$
- 2) $\frac{20!}{3!}$
- 3) ${}_{20}C_3$
- 4) ${}_{20}P_3$

208 What is a formula for the n th term of sequence B shown below?

$$B = 10, 12, 14, 16, \dots$$

- 1) $b_n = 8 + 2n$
- 2) $b_n = 10 + 2n$
- 3) $b_n = 10(2)^n$
- 4) $b_n = 10(2)^{n-1}$

209 If $f(x) = x^2 - 5$ and $g(x) = 6x$, then $g(f(x))$ is equal to

- 1) $6x^3 - 30x$
- 2) $6x^2 - 30$
- 3) $36x^2 - 5$
- 4) $x^2 + 6x - 5$

210 Four points on the graph of the function $f(x)$ are shown below.

$$\{(0, 1), (1, 2), (2, 4), (3, 8)\}$$

Which equation represents $f(x)$?

- 1) $f(x) = 2^x$
- 2) $f(x) = 2x$
- 3) $f(x) = x + 1$
- 4) $f(x) = \log_2 x$

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org


211 Which arithmetic sequence has a common difference of 4?

- 1) $\{0, 4n, 8n, 12n, \dots\}$
- 2) $\{n, 4n, 16n, 64n, \dots\}$
- 3) $\{n + 1, n + 5, n + 9, n + 13, \dots\}$
- 4) $\{n + 4, n + 16, n + 64, n + 256, \dots\}$

212 A survey completed at a large university asked 2,000 students to estimate the average number of hours they spend studying each week. Every tenth student entering the library was surveyed. The data showed that the mean number of hours that students spend studying was 15.7 per week. Which characteristic of the survey could create a bias in the results?

- 1) the size of the sample
- 2) the size of the population
- 3) the method of analyzing the data
- 4) the method of choosing the students who were surveyed

213 In the diagram below of right triangle KTW , $KW = 6$, $KT = 5$, and $m\angle KTW = 90$.



What is the measure of $\angle K$, to the nearest minute?

- 1) $33^\circ 33'$
- 2) $33^\circ 34'$
- 3) $33^\circ 55'$
- 4) $33^\circ 56'$

214 The fraction $\frac{3}{\sqrt{3a^2b}}$ is equivalent to

- 1) $\frac{1}{a\sqrt{b}}$
- 2) $\frac{\sqrt{b}}{ab}$
- 3) $\frac{\sqrt{3b}}{ab}$
- 4) $\frac{\sqrt{3}}{a}$

215 What is the fifteenth term of the sequence $5, -10, 20, -40, 80, \dots$?

- 1) $-163, 840$
- 2) $-81, 920$
- 3) $81, 920$
- 4) $327, 680$

216 Which task is *not* a component of an observational study?

- 1) The researcher decides who will make up the sample.
- 2) The researcher analyzes the data received from the sample.
- 3) The researcher gathers data from the sample, using surveys or taking measurements.
- 4) The researcher divides the sample into two groups, with one group acting as a control group.

- 217 The table below shows the first-quarter averages for Mr. Harper's statistics class.

Statistics Class Averages

Quarter Averages	Frequency
99	1
97	5
95	4
92	4
90	7
87	2
84	6
81	2
75	1
70	2
65	1

What is the population variance for this set of data?

- 1) 8.2
 - 2) 8.3
 - 3) 67.3
 - 4) 69.3
- 218 The expression $\frac{\sin^2 \theta + \cos^2 \theta}{1 - \sin^2 \theta}$ is equivalent to

- 1) $\cos^2 \theta$
- 2) $\sin^2 \theta$
- 3) $\sec^2 \theta$
- 4) $\csc^2 \theta$


- 219 The value of the expression $\sum_{r=3}^5 (-r^2 + r)$ is

- 1) -38
- 2) -12
- 3) 26
- 4) 62

- 220 The sides of a parallelogram measure 10 cm and 18 cm. One angle of the parallelogram measures 46 degrees. What is the area of the parallelogram, to the nearest square centimeter?

- 1) 65
- 2) 125
- 3) 129
- 4) 162

- 221 Which equation is represented by the graph below?



- 1) $y = 5^x$
- 2) $y = 0.5^x$
- 3) $y = 5^{-x}$
- 4) $y = 0.5^{-x}$

222 Written in simplest form, the expression $\frac{\frac{x}{4} - \frac{1}{x}}{\frac{1}{2x} + \frac{1}{4}}$

is equivalent to

- 1) $x - 1$
- 2) $x - 2$
- 3) $\frac{x - 2}{2}$
- 4) $\frac{x^2 - 4}{x + 2}$

223 Akeem invests \$25,000 in an account that pays 4.75% annual interest compounded continuously. Using the formula $A = Pe^{rt}$, where A = the amount in the account after t years, P = principal invested, and r = the annual interest rate, how many years, to the nearest tenth, will it take for Akeem's investment to triple?

- 1) 10.0
- 2) 14.6
- 3) 23.1
- 4) 24.0


224 The solution set of the equation $\sqrt{x + 3} = 3 - x$ is

- 1) {1}
- 2) {0}
- 3) {1, 6}
- 4) {2, 3}


225 The roots of the equation $x^2 - 10x + 25 = 0$ are

- 1) imaginary
- 2) real and irrational
- 3) real, rational, and equal
- 4) real, rational, and unequal


226 Which graph best represents the inequality $y + 6 \geq x^2 - x$?




1)



2)



3)



4)

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

227 The expression $\frac{4}{5 - \sqrt{13}}$ is equivalent to

- 1) $\frac{4\sqrt{13}}{5\sqrt{13} - 13}$
- 2) $\frac{4(5 - \sqrt{13})}{38}$
- 3) $\frac{5 + \sqrt{13}}{3}$
- 4) $\frac{4(5 + \sqrt{13})}{38}$

228 The expression $\log_8 64$ is equivalent to

- 1) 8
- 2) 2
- 3) $\frac{1}{2}$
- 4) $\frac{1}{8}$

229 The expression $(x^2 - 1)^{-\frac{2}{3}}$ is equivalent to

- 1) $\sqrt[3]{(x^2 - 1)^2}$
- 2) $\frac{1}{\sqrt[3]{(x^2 - 1)^2}}$
- 3) $\sqrt{(x^2 - 1)^3}$
- 4) $\frac{1}{\sqrt{(x^2 - 1)^3}}$

230 When $\frac{3}{2}x^2 - \frac{1}{4}x - 4$ is subtracted from


$\frac{5}{2}x^2 - \frac{3}{4}x + 1$, the difference is

- 1) $-x^2 + \frac{1}{2}x - 5$
- 2) $x^2 - \frac{1}{2}x + 5$
- 3) $-x^2 - x - 3$
- 4) $x^2 - x - 3$

231 Which graph does *not* represent a function?

- 1)
- 2)
- 3)
- 4)

232 What are the domain and the range of the function shown in the graph below?



- 1) $\{x|x > -4\}; \{y|y > 2\}$
- 2) $\{x|x \geq -4\}; \{y|y \geq 2\}$
- 3) $\{x|x > 2\}; \{y|y > -4\}$
- 4) $\{x|x \geq 2\}; \{y|y \geq -4\}$

233 Factored completely, the expression

$$12x^4 + 10x^3 - 12x^2$$


is equivalent to

- 1) $x^2(4x + 6)(3x - 2)$
- 2) $2(2x^2 + 3x)(3x^2 - 2x)$
- 3) $2x^2(2x - 3)(3x + 2)$
- 4) $2x^2(2x + 3)(3x - 2)$


234 An auditorium has 21 rows of seats. The first row has 18 seats, and each succeeding row has two more seats than the previous row. How many seats are in the auditorium?

- 1) 540
- 2) 567
- 3) 760
- 4) 798


235 If a function is defined by the equation $f(x) = 4^x$, which graph represents the inverse of this function?




1)



2)



3)



4)

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

236 For which equation does the sum of the roots equal -3 and the product of the roots equal 2 ?

- 1) $x^2 + 2x - 3 = 0$
- 2) $x^2 - 3x + 2 = 0$
- 3) $2x^2 + 6x + 4 = 0$
- 4) $2x^2 - 6x + 4 = 0$

237 What is the principal value of $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$?

- 1) -30°
- 2) 60°
- 3) 150°
- 4) 240°

238 For which equation does the sum of the roots equal $\frac{3}{4}$ and the product of the roots equal -2 ?

- 1) $4x^2 - 8x + 3 = 0$
- 2) $4x^2 + 8x + 3 = 0$
- 3) $4x^2 - 3x - 8 = 0$
- 4) $4x^2 + 3x - 2 = 0$

239 Which values of x are solutions of the equation

$$x^3 + x^2 - 2x = 0?$$

- 1) $0, 1, 2$
- 2) $0, 1, -2$
- 3) $0, -1, 2$
- 4) $0, -1, -2$

240 Mrs. Hill asked her students to express the sum $1 + 3 + 5 + 7 + 9 + \dots + 39$ using sigma notation. Four different student answers were given. Which student answer is correct?

1) $\sum_{k=1}^{20} (2k-1)$

2) $\sum_{k=2}^{40} (k-1)$

3) $\sum_{k=-1}^{37} (k+2)$

4) $\sum_{k=1}^{39} (2k-1)$

241 If $r = \sqrt[3]{\frac{A^2B}{C}}$, then $\log r$ can be represented by

1) $\frac{1}{6} \log A + \frac{1}{3} \log B - \log C$

2) $3(\log A^2 + \log B - \log C)$

3) $\frac{1}{3} \log(A^2 + B) - C$

4) $\frac{2}{3} \log A + \frac{1}{3} \log B - \frac{1}{3} \log C$

242 When $x^{-1} - 1$ is divided by $x - 1$, the quotient is

1) -1

2) $-\frac{1}{x}$

3) $\frac{1}{x^2}$

4) $\frac{1}{(x-1)^2}$

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org


243 What is the solution of the equation $2\log_4(5x) = 3$?

- 1) 6.4
- 2) 2.56
- 3) $\frac{9}{5}$
- 4) $\frac{8}{5}$

244 The roots of the equation $9x^2 + 3x - 4 = 0$ are

- 1) imaginary
- 2) real, rational, and equal
- 3) real, rational, and unequal
- 4) real, irrational, and unequal

245 In $\triangle ABC$, $a = 15$, $b = 14$, and $c = 13$, as shown in the diagram below. What is the $m\angle C$, to the nearest degree?



- 1) 53
- 2) 59
- 3) 67
- 4) 127


246 How many distinct triangles can be formed if $m\angle A = 35$, $a = 10$, and $b = 13$?

- 1) 1
- 2) 2
- 3) 3
- 4) 0

247 What is the value of x in the equation $\log_5 x = 4$?

- 1) 1.16
- 2) 20
- 3) 625
- 4) 1,024


248 In the diagram below of a unit circle, the ordered pair $\left(-\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}\right)$ represents the point where the terminal side of θ intersects the unit circle.



What is $m\angle\theta$?

- 1) 45
- 2) 135
- 3) 225
- 4) 240

- 249 Samantha constructs the scatter plot below from a set of data.



Based on her scatter plot, which regression model would be most appropriate?

- 1) exponential
 - 2) linear
 - 3) logarithmic
 - 4) power
- 250 The expression $4ab\sqrt{2b} - 3a\sqrt{18b^3} + 7ab\sqrt{6b}$ is equivalent to
- 1) $2ab\sqrt{6b}$
 - 2) $16ab\sqrt{2b}$
 - 3) $-5ab + 7ab\sqrt{6b}$
 - 4) $-5ab\sqrt{2b} + 7ab\sqrt{6b}$
- 251 An amateur bowler calculated his bowling average for the season. If the data are normally distributed, about how many of his 50 games were within one standard deviation of the mean?
- 1) 14
 - 2) 17
 - 3) 34
 - 4) 48

- 252 The expression $\cos^2\theta - \cos 2\theta$ is equivalent to
- 1) $\sin^2\theta$
 - 2) $-\sin^2\theta$
 - 3) $\cos^2\theta + 1$
 - 4) $-\cos^2\theta - 1$

- 253 A doctor wants to test the effectiveness of a new drug on her patients. She separates her sample of patients into two groups and administers the drug to only one of these groups. She then compares the results. Which type of study *best* describes this situation?
- 1) census
 - 2) survey
 - 3) observation
 - 4) controlled experiment

- 254 The expression $\cos 4x \cos 3x + \sin 4x \sin 3x$ is equivalent to
- 1) $\sin x$
 - 2) $\sin 7x$
 - 3) $\cos x$
 - 4) $\cos 7x$

- 255 What is the period of the function $f(\theta) = -2\cos 3\theta$?
- 1) π
 - 2) $\frac{2\pi}{3}$
 - 3) $\frac{3\pi}{2}$
 - 4) 2π

256 Three marbles are to be drawn at random, without replacement, from a bag containing 15 red marbles, 10 blue marbles, and 5 white marbles. Which expression can be used to calculate the probability of drawing 2 red marbles and 1 white marble from the bag?

- 1) $\frac{{}^{15}C_2 \cdot {}^5C_1}{{}^{30}C_3}$
- 2) $\frac{{}^{15}P_2 \cdot {}^5P_1}{{}^{30}C_3}$
- 3) $\frac{{}^{15}C_2 \cdot {}^5C_1}{{}^{30}P_3}$
- 4) $\frac{{}^{15}P_2 \cdot {}^5P_1}{{}^{30}P_3}$

257 In $\triangle ABC$, $a = 3$, $b = 5$, and $c = 7$. What is $m\angle C$?

- 1) 22
- 2) 38
- 3) 60
- 4) 120

258 The expression $\log_5\left(\frac{1}{25}\right)$ is equivalent to

- 1) $\frac{1}{2}$
- 2) 2
- 3) $-\frac{1}{2}$
- 4) -2

259 The product of $(3 + \sqrt{5})$ and $(3 - \sqrt{5})$ is

- 1) $4 - 6\sqrt{5}$
- 2) $14 - 6\sqrt{5}$
- 3) 14
- 4) 4


260 The number of minutes students took to complete a quiz is summarized in the table below.

Minutes	14	15	16	17	18	19	20
Number of Students	5	3	x	5	2	10	1

If the mean number of minutes was 17, which equation could be used to calculate the value of x ?

- 1) $17 = \frac{119 + x}{x}$
- 2) $17 = \frac{119 + 16x}{x}$
- 3) $17 = \frac{446 + x}{26 + x}$
- 4) $17 = \frac{446 + 16x}{26 + x}$

261 Which equation represents the circle shown in the graph below that passes through the point $(0, -1)$?



- 1) $(x - 3)^2 + (y + 4)^2 = 16$
- 2) $(x - 3)^2 + (y + 4)^2 = 18$
- 3) $(x + 3)^2 + (y - 4)^2 = 16$
- 4) $(x + 3)^2 + (y - 4)^2 = 18$

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

262 What is the formula for the n th term of the sequence 54, 18, 6, ...?

- 1) $a_n = 6\left(\frac{1}{3}\right)^n$
- 2) $a_n = 6\left(\frac{1}{3}\right)^{n-1}$
- 3) $a_n = 54\left(\frac{1}{3}\right)^n$
- 4) $a_n = 54\left(\frac{1}{3}\right)^{n-1}$

263 Which two functions are inverse functions of each other?

- 1) $f(x) = \sin x$ and $g(x) = \cos(x)$
- 2) $f(x) = 3 + 8x$ and $g(x) = 3 - 8x$
- 3) $f(x) = e^x$ and $g(x) = \ln x$
- 4) $f(x) = 2x - 4$ and $g(x) = -\frac{1}{2}x + 4$

264 The expression $\frac{2x+4}{\sqrt{x+2}}$ is equivalent to

- 1) $\frac{(2x+4)\sqrt{x-2}}{x-2}$
- 2) $\frac{(2x+4)\sqrt{x-2}}{x-4}$
- 3) $2\sqrt{x-2}$
- 4) $2\sqrt{x+2}$

265 What is the number of degrees in an angle whose radian measure is $\frac{11\pi}{12}$?

- 1) 150
- 2) 165
- 3) 330
- 4) 518

266 If $f(x) = \frac{1}{2}x - 3$ and $g(x) = 2x + 5$, what is the value of $(g \circ f)(4)$?

- 1) -13
- 2) 3.5
- 3) 3
- 4) 6

267 What is the domain of the function

$$f(x) = \sqrt{x-2} + 3?$$


- 1) $(-\infty, \infty)$
- 2) $(2, \infty)$
- 3) $[2, \infty)$
- 4) $[3, \infty)$

268 What is the solution set of the equation

$$|4a + 6| - 4a = -10?$$

- 1) \emptyset
- 2) $\{0\}$
- 3) $\left\{\frac{1}{2}\right\}$
- 4) $\left\{0, \frac{1}{2}\right\}$

- 269 A dartboard is shown in the diagram below. The two lines intersect at the center of the circle, and the central angle in sector 2 measures $\frac{2\pi}{3}$.



If darts thrown at this board are equally likely to land anywhere on the board, what is the probability that a dart that hits the board will land in either sector 1 or sector 3?

- 1) $\frac{1}{6}$
 - 2) $\frac{1}{3}$
 - 3) $\frac{1}{2}$
 - 4) $\frac{2}{3}$
- 270 What is the common ratio of the geometric sequence whose first term is 27 and fourth term is 64?
- 1) $\frac{3}{4}$
 - 2) $\frac{64}{81}$
 - 3) $\frac{4}{3}$
 - 4) $\frac{37}{3}$


- 271 Expressed as a function of a positive acute angle, $\cos(-305^\circ)$ is equal to

- 1) $-\cos 55^\circ$
- 2) $\cos 55^\circ$
- 3) $-\sin 55^\circ$
- 4) $\sin 55^\circ$

- 272 In $\triangle ABC$, $m\angle A = 120$, $b = 10$, and $c = 18$. What is the area of $\triangle ABC$ to the nearest square inch?

- 1) 52
- 2) 78
- 3) 90
- 4) 156

- 273 Which equation is represented by the graph below?



- 1) $y = \cot x$
- 2) $y = \csc x$
- 3) $y = \sec x$
- 4) $y = \tan x$

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

274 What is the fourth term in the expansion of

$$(3x - 2)^5?$$

- 1) $-720x^2$
- 2) $-240x$
- 3) $720x^2$
- 4) $1,080x^3$

275 If $a = 3$ and $b = -2$, what is the value of the

$$\text{expression } \frac{a^{-2}}{b^{-3}}?$$

- 1) $-\frac{9}{8}$
- 2) -1
- 3) $-\frac{8}{9}$
- 4) $\frac{8}{9}$

276 Ms. Bell's mathematics class consists of 4 sophomores, 10 juniors, and 5 seniors. How many different ways can Ms. Bell create a four-member committee of juniors if each junior has an equal chance of being selected?

- 1) 210
- 2) 3,876
- 3) 5,040
- 4) 93,024

277 Which function is *not* one-to-one?

- 1) $\{(0, 1), (1, 2), (2, 3), (3, 4)\}$
- 2) $\{(0, 0), (1, 1), (2, 2), (3, 3)\}$
- 3) $\{(0, 1), (1, 0), (2, 3), (3, 2)\}$
- 4) $\{(0, 1), (1, 0), (2, 0), (3, 2)\}$

278 What is the radian measure of an angle whose measure is -420° ?

- 1) $-\frac{7\pi}{3}$
- 2) $-\frac{7\pi}{6}$
- 3) $\frac{7\pi}{6}$
- 4) $\frac{7\pi}{3}$

279 The value of $\tan 126^\circ 43'$ to the *nearest ten-thousandth* is

- 1) -1.3407
- 2) -1.3408
- 3) -1.3548
- 4) -1.3549

280 Which expression is equivalent to $\frac{\sqrt{3} + 5}{\sqrt{3} - 5}$?

- 1) $-\frac{14 + 5\sqrt{3}}{11}$
- 2) $-\frac{17 + 5\sqrt{3}}{11}$
- 3) $\frac{14 + 5\sqrt{3}}{14}$
- 4) $\frac{17 + 5\sqrt{3}}{14}$

281 What is the coefficient of the fourth term in the expansion of $(a - 4b)^9$?

- 1) $-5,376$
- 2) -336
- 3) 336
- 4) $5,376$

282 Which formula can be used to determine the total number of different eight-letter arrangements that can be formed using the letters in the word *DEADLINE*?

- 1) $8!$
- 2) $\frac{8!}{4!}$
- 3) $\frac{8!}{2! + 2!}$
- 4) $\frac{8!}{2! \cdot 2!}$





283 The expression $\sqrt[4]{16x^2y^7}$ is equivalent to

- 1) $2x^{\frac{1}{2}}y^{\frac{7}{4}}$
- 2) $2x^8y^{28}$
- 3) $4x^{\frac{1}{2}}y^{\frac{7}{4}}$
- 4) $4x^8y^{28}$

284 If $\angle A$ is acute and $\tan A = \frac{2}{3}$, then

- 1) $\cot A = \frac{2}{3}$
- 2) $\cot A = \frac{1}{3}$
- 3) $\cot(90^\circ - A) = \frac{2}{3}$
- 4) $\cot(90^\circ - A) = \frac{1}{3}$

285 Which graph represents a one-to-one function?

- 1) 
- 2) 
- 3) 
- 4) 

286 The value of the expression $2 \sum_{n=0}^2 (n^2 + 2^n)$ is

- 1) 12
- 2) 22
- 3) 24
- 4) 26





Algebra 2/Trigonometry Multiple Choice Regents Exam Questions

www.jmap.org

287 The equation $x^2 + y^2 - 2x + 6y + 3 = 0$ is equivalent to

- 1) $(x - 1)^2 + (y + 3)^2 = -3$
- 2) $(x - 1)^2 + (y + 3)^2 = 7$
- 3) $(x + 1)^2 + (y + 3)^2 = 7$
- 4) $(x + 1)^2 + (y + 3)^2 = 10$

288 Which graph does *not* represent a function?

- 1) 
- 2) 
- 3) 
- 4) 

289 The expression $2\log x - (3\log y + \log z)$ is equivalent to

- 1) $\log \frac{x^2}{y^3 z}$
- 2) $\log \frac{x^2 z}{y^3}$
- 3) $\log \frac{2x}{3yz}$
- 4) $\log \frac{2xz}{3y}$


290 Brian correctly used a method of completing the square to solve the equation $x^2 + 7x - 11 = 0$. Brian's first step was to rewrite the equation as $x^2 + 7x = 11$. He then added a number to both sides of the equation. Which number did he add?

- 1) $\frac{7}{2}$
- 2) $\frac{49}{4}$
- 3) $\frac{49}{2}$
- 4) 49


291 The principal would like to assemble a committee of 8 students from the 15-member student council. How many different committees can be chosen?

- 1) 120
- 2) 6,435
- 3) 32,432,400
- 4) 259,459,200


292 Which graph represents the equation $y = \cos^{-1}x$?




1)



2)



3)



4)

293 What is the common difference of the arithmetic sequence 5, 8, 11, 14?

- 1) $\frac{8}{5}$
- 2) -3
- 3) 3
- 4) 9

294 The function $f(x) = \tan x$ is defined in such a way that $f^{-1}(x)$ is a function. What can be the domain of $f(x)$?

- 1) $\{x \mid 0 \leq x \leq \pi\}$
- 2) $\{x \mid 0 \leq x \leq 2\pi\}$
- 3) $\left\{x \mid -\frac{\pi}{2} < x < \frac{\pi}{2}\right\}$
- 4) $\left\{x \mid -\frac{\pi}{2} < x < \frac{3\pi}{2}\right\}$

295 If $\sin^{-1}\left(\frac{5}{8}\right) = A$, then

- 1) $\sin A = \frac{5}{8}$
- 2) $\sin A = \frac{8}{5}$
- 3) $\cos A = \frac{5}{8}$
- 4) $\cos A = \frac{8}{5}$

296 Factored completely, the expression $6x - x^3 - x^2$ is equivalent to

- 1) $x(x+3)(x-2)$
- 2) $x(x-3)(x+2)$
- 3) $-x(x-3)(x+2)$
- 4) $-x(x+3)(x-2)$

297 Which value of r represents data with a strong negative linear correlation between two variables?


- 1) -1.07
- 2) -0.89
- 3) -0.14
- 4) 0.92

Algebra 2/Trigonometry 2 Point Regents Exam Questions

298 Show that $\sec \theta \sin \theta \cot \theta = 1$ is an identity.

299 Express $\cos \theta (\sec \theta - \cos \theta)$, in terms of $\sin \theta$.

300 Determine the solution of the inequality $|3 - 2x| \geq 7$. [The use of the grid below is optional.]



301 On a multiple-choice test, Abby randomly guesses on all seven questions. Each question has four choices. Find the probability, to the *nearest thousandth*, that Abby gets *exactly* three questions correct.

302 Find, algebraically, the measure of the obtuse angle, to the *nearest degree*, that satisfies the equation $5 \csc \theta = 8$.

303 Write a quadratic equation such that the sum of its roots is 6 and the product of its roots is -27 .

304 Starting with $\sin^2 A + \cos^2 A = 1$, derive the formula $\tan^2 A + 1 = \sec^2 A$.

305 A cup of soup is left on a countertop to cool. The table below gives the temperatures, in degrees Fahrenheit, of the soup recorded over a 10-minute period.


Time in Minutes (x)	Temperature in $^{\circ}\text{F}$ (y)
0	180.2
2	165.8
4	146.3
6	135.4
8	127.7
10	110.5

Write an exponential regression equation for the data, rounding all values to the *nearest thousandth*.

306 Convert 3 radians to degrees and express the answer to the *nearest minute*.

307 Convert 2.5 radians to degrees, and express the answer to the *nearest minute*.

308 The graph below represents the function $y = f(x)$.



State the domain and range of this function.

309 If θ is an angle in standard position and its terminal side passes through the point $(-3, 2)$, find the exact value of $\csc \theta$.

310 Howard collected fish eggs from a pond behind his house so he could determine whether sunlight had an effect on how many of the eggs hatched. After he collected the eggs, he divided them into two tanks. He put both tanks outside near the pond, and he covered one of the tanks with a box to block out all sunlight. State whether Howard's investigation was an example of a controlled experiment, an observation, or a survey. Justify your response.

311 The table below shows the number of new stores in a coffee shop chain that opened during the years 1986 through 1994.

Year	Number of New Stores
1986	14
1987	27
1988	48
1989	80
1990	110
1991	153
1992	261
1993	403
1994	681

Using $x = 1$ to represent the year 1986 and y to represent the number of new stores, write the exponential regression equation for these data. Round all values to the *nearest thousandth*.

312 Solve algebraically for x : $16^{2x+3} = 64^{x+2}$

313 If $f(x) = x^2 - 6$ and $g(x) = 2^x - 1$, determine the value of $(g \circ f)(-3)$.

314 Solve algebraically for x : $4 - \sqrt{2x - 5} = 1$

315 Factor the expression $12t^8 - 75t^4$ completely.

Algebra 2/Trigonometry 2 Point Regents Exam Questions

www.jmap.org

316 Find the third term in the recursive sequence
 $a_{k+1} = 2a_k - 1$, where $a_1 = 3$.

317 Solve algebraically for x : $\log_{27}(2x - 1) = \frac{4}{3}$

318 Find, to the *nearest tenth of a degree*, the angle whose measure is 2.5 radians.

319 Determine the sum and the product of the roots of $3x^2 = 11x - 6$.

320 The table below shows the concentration of ozone in Earth's atmosphere at different altitudes. Write the exponential regression equation that models these data, rounding *all* values to the *nearest thousandth*.

Concentration of Ozone	
Altitude (x)	Ozone Units (y)
0	0.7
5	0.6
10	1.1
15	3.0
20	4.9


321 Solve for x : $\frac{4x}{x-3} = 2 + \frac{12}{x-3}$

322 Find the solution of the inequality $x^2 - 4x > 5$, algebraically.

323 The formula for continuously compounded interest is $A = Pe^{rt}$, where A is the amount of money in the account, P is the initial investment, r is the interest rate, and t is the time in years. Using the formula, determine, to the *nearest dollar*, the amount in the account after 8 years if \$750 is invested at an annual rate of 3%.

324 In an arithmetic sequence, $a_4 = 19$ and $a_7 = 31$. Determine a formula for a_n , the n^{th} term of this sequence.

325 Write an equation of the circle shown in the diagram below.



326 If $\sec(a + 15)^\circ = \csc(2a)^\circ$, find the smallest positive value of a , in degrees.

Algebra 2/Trigonometry 2 Point Regents Exam Questions


www.jmap.org

327 If $g(x) = (ax\sqrt{1-x})^2$, express $g(10)$ in simplest form.

328 Determine the value of n in simplest form:

$$i^{13} + i^{18} + i^{31} + n = 0$$

329 The graph of the equation $y = \left(\frac{1}{2}\right)^x$ has an asymptote. On the grid below, sketch the graph of $y = \left(\frac{1}{2}\right)^x$ and write the equation of this asymptote.



330 A blood bank needs twenty people to help with a blood drive. Twenty-five people have volunteered. Find how many different groups of twenty can be formed from the twenty-five volunteers.

331 Solve $\sec x - \sqrt{2} = 0$ algebraically for all values of x in $0^\circ \leq x < 360^\circ$.

332 Express the exact value of $\csc 60^\circ$, with a rational denominator.

333 Express in simplest form: $\frac{\frac{1}{2} - \frac{4}{d}}{\frac{1}{d} + \frac{3}{2d}}$

334 Find the sum and product of the roots of the equation $5x^2 + 11x - 3 = 0$.

335 Express $4xi + 5yi^8 + 6xi^3 + 2yi^4$ in simplest $a + bi$ form.


336 A committee of 5 members is to be randomly selected from a group of 9 teachers and 20 students. Determine how many different committees can be formed if 2 members must be teachers and 3 members must be students.

337 Determine algebraically the x -coordinate of all points where the graphs of $xy = 10$ and $y = x + 3$ intersect.

Algebra 2/Trigonometry 2 Point Regents Exam Questions

www.jmap.org

- 338 Write an equation for the graph of the trigonometric function shown below.



- 339 Solve $|-4x + 5| < 13$ algebraically for x .

- 340 Evaluate $e^{x \ln y}$ when $x = 3$ and $y = 2$.


- 341 Express $5\sqrt{3x^3} - 2\sqrt{27x^3}$ in simplest radical form.

- 342 Express in simplest form: $\sqrt[3]{\frac{a^6 b^9}{-64}}$

- 343 Express $\frac{5}{3 - \sqrt{2}}$ with a rational denominator, in simplest radical form.

- 344 Use the discriminant to determine all values of k that would result in the equation $x^2 - kx + 4 = 0$ having equal roots.

- 345 On the axes below, for $-2 \leq x \leq 2$, graph $y = 2^{x+1} - 3$.




- 346 Find the total number of different twelve-letter arrangements that can be formed using the letters in the word *PENNSYLVANIA*.

- 347 Find, to the *nearest tenth of a square foot*, the area of a rhombus that has a side of 6 feet and an angle of 50° .

348 Factor completely: $10ax^2 - 23ax - 5a$

349 Determine the sum and the product of the roots of the equation $12x^2 + x - 6 = 0$.

350 On the unit circle shown in the diagram below, sketch an angle, in standard position, whose degree measure is 240 and find the exact value of $\sin 240^\circ$.



351 Find, to the *nearest minute*, the angle whose measure is 3.45 radians.

352 Express the product of $\cos 30^\circ$ and $\sin 45^\circ$ in simplest radical form.


353 Find the first four terms of the recursive sequence defined below.

$$a_1 = -3$$

$$a_n = a_{(n-1)} - n$$

354 Solve the equation $2 \tan C - 3 = 3 \tan C - 4$ algebraically for all values of C in the interval $0^\circ \leq C < 360^\circ$.

355 A circle shown in the diagram below has a center of $(-5, 3)$ and passes through point $(-1, 7)$.




Write an equation that represents the circle.


356 In a certain school, the heights of the population of girls are normally distributed, with a mean of 63 inches and a standard deviation of 2 inches. If there are 450 girls in the school, determine how many of the girls are *shorter than* 60 inches. Round the answer to the *nearest integer*.

357 Solve algebraically for x : $\log_{5x-1} 4 = \frac{1}{3}$

358 Write an equation of the circle shown in the graph below.



359 Circle O shown below has a radius of 12 centimeters. To the *nearest tenth of a centimeter*, determine the length of the arc, x , subtended by an angle of $83^\circ 50'$.



360 Solve the equation $6x^2 - 2x - 3 = 0$ and express the answer in simplest radical form.

361 The heights, in inches, of 10 high school varsity basketball players are 78, 79, 79, 72, 75, 71, 74, 74, 83, and 71. Find the interquartile range of this data set.

362 Express the sum $7 + 14 + 21 + 28 + \dots + 105$ using sigma notation.

363 For a given set of rectangles, the length is inversely proportional to the width. In one of these rectangles, the length is 12 and the width is 6. For this set of rectangles, calculate the width of a rectangle whose length is 9.

364 Determine the sum of the first twenty terms of the sequence whose first five terms are 5, 14, 23, 32, 41.

365 Express $\frac{\sqrt{108x^5y^8}}{\sqrt{6xy^5}}$ in simplest radical form.

366 Evaluate: $10 + \sum_{n=1}^5 (n^3 - 1)$

367 In a study of 82 video game players, the researchers found that the ages of these players were normally distributed, with a mean age of 17 years and a standard deviation of 3 years. Determine if there were 15 video game players in this study over the age of 20. Justify your answer.

368 Multiply $x + yi$ by its conjugate, and express the product in simplest form.

369 The scores of one class on the Unit 2 mathematics test are shown in the table below.


Unit 2 Mathematics Test

Test Score	Frequency
96	1
92	2
84	5
80	3
76	6
72	3
68	2

Find the population standard deviation of these scores, to the *nearest tenth*.

370 Express $\frac{\cot x \sin x}{\sec x}$ as a single trigonometric function, in simplest form, for all values of x for which it is defined.

371 The function $f(x)$ is graphed on the set of axes below. On the same set of axes, graph $f(x + 1) + 2$.



372 Assume that the ages of first-year college students are normally distributed with a mean of 19 years and standard deviation of 1 year. To the *nearest integer*, find the percentage of first-year college students who are between the ages of 18 years and 20 years, inclusive. To the *nearest integer*, find the percentage of first-year college students who are 20 years old or older.

373 Solve $|2x - 3| > 5$ algebraically.

374 Simplify the expression $\frac{3x^{-4}y^5}{(2x^3y^{-7})^{-2}}$ and write the answer using only positive exponents.

375 If $f(x) = x^2 - 6$, find $f^{-1}(x)$.

376 Find, to the *nearest tenth*, the radian measure of 216° .

377 The number of bacteria present in a Petri dish can be modeled by the function $N = 50e^{3t}$, where N is the number of bacteria present in the Petri dish after t hours. Using this model, determine, to the *nearest hundredth*, the number of hours it will take for N to reach 30,700.

378 Express $\left(\frac{2}{3}x - 1\right)^2$ as a trinomial.

379 The following is a list of the individual points scored by all twelve members of the Webster High School basketball team at a recent game:
2 2 3 4 6 7 9 10 10 11 12 14
Find the interquartile range for this set of data.

380 Express the product of $\left(\frac{1}{2}y^2 - \frac{1}{3}y\right)$ and $\left(12y + \frac{3}{5}\right)$ as a trinomial.

381 Evaluate: $\sum_{n=1}^3 (-n^4 - n)$

382 The two sides and included angle of a parallelogram are 18, 22, and 60° . Find its exact area in simplest form.

383 Matt places \$1,200 in an investment account earning an annual rate of 6.5%, compounded continuously. Using the formula $V = Pe^{rt}$, where V is the value of the account in t years, P is the principal initially invested, e is the base of a natural logarithm, and r is the rate of interest, determine the amount of money, to the *nearest cent*, that Matt will have in the account after 10 years.

384 Two sides of a parallelogram are 24 feet and 30 feet. The measure of the angle between these sides is 57° . Find the area of the parallelogram, to the *nearest square foot*.

385 Find the number of possible different 10-letter arrangements using the letters of the word "STATISTICS."

Algebra 2/Trigonometry 4 Point Regents Exam Questions

386 Solve algebraically for all values of x :
 $\log_{(x+4)}(17x - 4) = 2$

387 The probability that a professional baseball player will get a hit is $\frac{1}{3}$. Calculate the exact probability that he will get *at least* 3 hits in 5 attempts.

388 Graph the inequality $-3|6 - x| < -15$ for x . Graph the solution on the line below.



389 The data collected by a biologist showing the growth of a colony of bacteria at the end of each hour are displayed in the table below.

Time, hour, (x)	Population (y)
0	250
1	330
2	580
3	800
4	1650
5	3000

Write an exponential regression equation to model these data. Round all values to the *nearest thousandth*. Assuming this trend continues, use this equation to estimate, to the nearest *ten*, the number of bacteria in the colony at the end of 7 hours.

390 A population of single-celled organisms was grown in a Petri dish over a period of 16 hours. The number of organisms at a given time is recorded in the table below.

Time, hrs (x)	Number of Organisms (y)
0	25
2	36
4	52
6	68
8	85
10	104
12	142
16	260


Determine the exponential regression equation model for these data, rounding all values to the *nearest ten-thousandth*. Using this equation, predict the number of single-celled organisms, to the *nearest whole number*, at the end of the 18th hour.

391 The probability that the Stormville Sluggers will win a baseball game is $\frac{2}{3}$. Determine the probability, to the *nearest thousandth*, that the Stormville Sluggers will win *at least* 6 of their next 8 games.

392 Solve the equation $8x^3 + 4x^2 - 18x - 9 = 0$ algebraically for all values of x .

393 The measures of the angles between the resultant and two applied forces are 60° and 45° , and the magnitude of the resultant is 27 pounds. Find, to the *nearest pound*, the magnitude of each applied force.

394 As shown in the diagram below, fire-tracking station A is 100 miles due west of fire-tracking station B . A forest fire is spotted at F , on a bearing 47° northeast of station A and 15° northeast of station B . Determine, to the *nearest tenth of a mile*, the distance the fire is from *both* station A and station B . [N represents due north.]



395 Express as a single fraction the exact value of $\sin 75^\circ$.

396 The table below shows the results of an experiment involving the growth of bacteria.

Time (x) (in minutes)	1	3	5	7	9	11
Number of Bacteria (y)	2	25	81	175	310	497

Write a power regression equation for this set of data, rounding all values to *three decimal places*. Using this equation, predict the bacteria's growth, to the *nearest integer*, after 15 minutes.

397 The letters of any word can be rearranged. Carol believes that the number of different 9-letter arrangements of the word "TENNESSEE" is greater than the number of different 7-letter arrangements of the word "VERMONT." Is she correct? Justify your answer.

398 Find all values of θ in the interval $0^\circ \leq \theta < 360^\circ$ that satisfy the equation $\sin 2\theta = \sin \theta$.

399 A study shows that 35% of the fish caught in a local lake had high levels of mercury. Suppose that 10 fish were caught from this lake. Find, to the *nearest tenth of a percent*, the probability that *at least* 8 of the 10 fish caught did *not* contain high levels of mercury.

400 A ranch in the Australian Outback is shaped like triangle ACE , with $m\angle A = 42$, $m\angle E = 103$, and $AC = 15$ miles. Find the area of the ranch, to the *nearest square mile*.

401 The members of a men's club have a choice of wearing black or red vests to their club meetings. A study done over a period of many years determined that the percentage of black vests worn is 60%. If there are 10 men at a club meeting on a given night, what is the probability, to the *nearest thousandth*, that *at least* 8 of the vests worn will be black?

Algebra 2/Trigonometry 4 Point Regents Exam Questions


www.jmap.org

- 402 During a particular month, a local company surveyed all its employees to determine their travel times to work, in minutes. The data for all 15 employees are shown below.

25 55 40 65 29
45 59 35 25 37
52 30 8 40 55

Determine the number of employees whose travel time is within one standard deviation of the mean.

- 403 The diagram below shows the plans for a cell phone tower. A guy wire attached to the top of the tower makes an angle of 65 degrees with the ground. From a point on the ground 100 feet from the end of the guy wire, the angle of elevation to the top of the tower is 32 degrees. Find the height of the tower, to the *nearest foot*.



- 404 Two sides of a parallelogram measure 27 cm and 32 cm. The included angle measures 48° . Find the length of the longer diagonal of the parallelogram, to the *nearest centimeter*.
- 405 Solve algebraically for all exact values of x in the interval $0 \leq x < 2\pi$: $2\sin^2 x + 5\sin x = 3$

- 406 Solve the equation below algebraically, and express the result in simplest radical form:

$$\frac{13}{x} = 10 - x$$

- 407 Write the binomial expansion of $(2x - 1)^5$ as a polynomial in simplest form.

- 408 Express in simplest terms: $\frac{1 + \frac{3}{x}}{1 - \frac{5}{x} - \frac{24}{x^2}}$

- 409 Because Sam's backyard gets very little sunlight, the probability that a geranium planted there will flower is 0.28. Sam planted five geraniums. Determine the probability, to the *nearest thousandth*, that *at least* four geraniums will flower.

- 410 If $\tan A = \frac{2}{3}$ and $\sin B = \frac{5}{\sqrt{41}}$ and angles A and B are in Quadrant I, find the value of $\tan(A + B)$.

- 411 If $\log_4 x = 2.5$ and $\log_y 125 = -\frac{3}{2}$, find the numerical value of $\frac{x}{y}$, in simplest form.

412 Solve $x^3 + 5x^2 = 4x + 20$ algebraically.

413 Whenever Sara rents a movie, the probability that it is a horror movie is 0.57. Of the next five movies she rents, determine the probability, to the *nearest hundredth*, that *no more than* two of these rentals are horror movies.

414 Express in simplest form: $\frac{\frac{4 - x^2}{x^2 + 7x + 12}}{\frac{2x - 4}{x + 3}}$

415 Solve $2x^2 - 12x + 4 = 0$ by completing the square, expressing the result in simplest radical form.

416 Ten teams competed in a cheerleading competition at a local high school. Their scores were 29, 28, 39, 37, 45, 40, 41, 38, 37, and 48. How many scores are within one population standard deviation from the mean? For these data, what is the interquartile range?

417 Solve algebraically for x : $\frac{1}{x + 3} - \frac{2}{3 - x} = \frac{4}{x^2 - 9}$

418 In $\triangle ABC$, $m\angle A = 32$, $a = 12$, and $b = 10$. Find the measures of the missing angles and side of $\triangle ABC$. Round each measure to the *nearest tenth*.

Algebra 2/Trigonometry 6 Point Regents Exam Questions

- 419 Solve algebraically for all values of x :

$$81^{x^3 + 2x^2} = 27^{\frac{5x}{3}}$$

- 420 Solve algebraically for x :

$$\sqrt{x^2 + x - 1} + 11x = 7x + 3$$

- 421 Perform the indicated operations and simplify completely:

$$\frac{x^3 - 3x^2 + 6x - 18}{x^2 - 4x} \cdot \frac{2x - 4}{x^4 - 3x^3} \div \frac{x^2 + 2x - 8}{16 - x^2}$$

- 422 Solve algebraically for x : $\log_{x+3} \frac{x^3 + x - 2}{x} = 2$

- 423 Solve algebraically for all values of x :

$$x^4 + 4x^3 + 4x^2 = -16x$$

- 424 Solve the following systems of equations algebraically: $5 = y - x$

$$4x^2 = -17x + y + 4$$

- 425 Solve algebraically for all values of x :

$$\log_{(x+3)}(2x+3) + \log_{(x+3)}(x+5) = 2$$

- 426 In a triangle, two sides that measure 6 cm and 10 cm form an angle that measures 80° . Find, to the *nearest degree*, the measure of the smallest angle in the triangle.

- 427 The temperature, T , of a given cup of hot chocolate after it has been cooling for t minutes can best be modeled by the function below, where T_0 is the temperature of the room and k is a constant.

$$\ln(T - T_0) = -kt + 4.718$$

A cup of hot chocolate is placed in a room that has a temperature of 68° . After 3 minutes, the temperature of the hot chocolate is 150° . Compute the value of k to the nearest thousandth. [Only an algebraic solution can receive full credit.] Using this value of k , find the temperature, T , of this cup of hot chocolate if it has been sitting in this room for a total of 10 minutes. Express your answer to the *nearest degree*. [Only an algebraic solution can receive full credit.]

- 428 Two forces of 25 newtons and 85 newtons acting on a body form an angle of 55° . Find the magnitude of the resultant force, to the *nearest hundredth of a newton*. Find the measure, to the *nearest degree*, of the angle formed between the resultant and the larger force.

- 429 Two forces of 40 pounds and 28 pounds act on an object. The angle between the two forces is 65° . Find the magnitude of the resultant force, to the *nearest pound*. Using this answer, find the measure of the angle formed between the resultant and the *smaller* force, to the *nearest degree*.

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions Answer Section

1 ANS: 3

$$(-5)^2 - 4(2)(0) = 25$$

PTS: 2 REF: 061423a2 STA: A2.A.2 TOP: Using the Discriminant
KEY: determine equation given nature of roots

2 ANS: 2 PTS: 2 REF: 011222a2 STA: A2.A.39
TOP: Domain and Range KEY: real domain

3 ANS: 4 PTS: 2 REF: 011409a2 STA: A2.S.10
TOP: Permutations

4 ANS: 1

$$\frac{1}{2}(7.4)(3.8)\sin 126 \approx 11.4$$

PTS: 2 REF: 011218a2 STA: A2.A.74 TOP: Using Trigonometry to Find Area
KEY: basic

5 ANS: 4 PTS: 2 REF: 061206a2 STA: A2.A.60
TOP: Unit Circle

6 ANS: 4

$$\sin(\theta + 90) = \sin \theta \cdot \cos 90 + \cos \theta \cdot \sin 90 = \sin \theta \cdot (0) + \cos \theta \cdot (1) = \cos \theta$$

PTS: 2 REF: 061309a2 STA: A2.A.76 TOP: Angle Sum and Difference Identities
KEY: identities

7 ANS: 4

$$r = \sqrt{(6-3)^2 + (5-(-4))^2} = \sqrt{9+81} = \sqrt{90}$$

PTS: 2 REF: 061415a2 STA: A2.A.48 TOP: Equations of Circles

8 ANS: 3

$6n^{-1} < 4n^{-1}$. Flip sign when multiplying each side of the inequality by n , since a negative number.

$$\frac{6}{n} < \frac{4}{n}$$

$$6 > 4$$

PTS: 2 REF: 061314a2 STA: A2.N.1 TOP: Negative and Fractional Exponents

9 ANS: 4 PTS: 2 REF: 011323a2 STA: A2.A.2
TOP: Using the Discriminant KEY: determine nature of roots given equation

10 ANS: 2

$$(x+2)^2 = -9$$

$$x+2 = \pm\sqrt{-9}$$

$$x = -2 \pm 3i$$

PTS: 2 REF: 011408a2 STA: A2.A.24 TOP: Completing the Square

11 ANS: 2

$$\text{Top } 6.7\% = 1.5 \text{ s.d.} + \sigma = 1.5(104) + 576 = 732$$

PTS: 2

REF: 011420a2

STA: A2.S.5

TOP: Normal Distributions

KEY: predict

12 ANS: 2

$$60 = -16t^2 + 5t + 105 \quad t = \frac{-5 \pm \sqrt{5^2 - 4(-16)(45)}}{2(-16)} \approx \frac{-5 \pm 53.89}{-32} \approx 1.84$$

$$0 = -16t^2 + 5t + 45$$

PTS: 2

REF: 061424a2

STA: A2.A.25

TOP: Quadratics with Irrational Solutions

13 ANS: 2

PTS: 2

REF: 011301a2

STA: A2.A.53

TOP: Graphing Exponential Functions

14 ANS: 3

PTS: 2

REF: 061306a2

STA: A2.A.72

TOP: Identifying the Equation of a Trigonometric Graph

15 ANS: 2

$$b^2 - 4ac = (-9)^2 - 4(2)(4) = 81 - 32 = 49$$

PTS: 2

REF: 011411a2

STA: A2.A.2

TOP: Using the Discriminant

KEY: determine nature of roots given equation

16 ANS: 2

$$x^3 + 3x^2 - 4x - 12$$

$$x^2(x+3) - 4(x+3)$$

$$(x^2 - 4)(x+3)$$

$$(x+2)(x-2)(x+3)$$

PTS: 2

REF: 061214a2

STA: A2.A.7

TOP: Factoring by Grouping

17 ANS: 2

PTS: 2

REF: 061301a2

STA: A2.S.1

TOP: Analysis of Data

18 ANS: 2

$$\frac{8\pi}{5} \cdot \frac{180}{\pi} = 288$$

PTS: 2

REF: 061302a2

STA: A2.M.2

TOP: Radian Measure

KEY: degrees

19 ANS: 4

$$\frac{10}{4} = 2.5$$

PTS: 2

REF: 011217a2

STA: A2.A.29

TOP: Sequences

20 ANS: 3

$$\frac{a + \frac{b}{c}}{d - \frac{b}{c}} = \frac{\frac{ac + b}{c}}{\frac{cd - b}{c}} = \frac{ac + b}{c} \cdot \frac{c}{cd - b} = \frac{ac + b}{cd - b}$$

PTS: 2 REF: 011405a2 STA: A2.A.17 TOP: Complex Fractions

21 ANS: 3

$$s = \theta r = \frac{2\pi}{8} \cdot 6 = \frac{3\pi}{2}$$

PTS: 2 REF: 061212a2 STA: A2.A.61 TOP: Arc Length

KEY: arc length

22 ANS: 1

$$5x + 29 = (x + 3)^2 \quad . \quad (-5) + 3 \text{ shows an extraneous solution.}$$

$$5x + 29 = x^2 + 6x + 9$$

$$0 = x^2 + x - 20$$

$$0 = (x + 5)(x - 4)$$

$$x = -5, 4$$

PTS: 2 REF: 061213a2 STA: A2.A.22 TOP: Solving Radicals

KEY: extraneous solutions

23 ANS: 3

$${}_3C_1 \cdot {}_5C_2 = 3 \cdot 10 = 30$$

PTS: 2 REF: 061422a2 STA: A2.S.12 TOP: Combinations

24 ANS: 2

$$\log x^2 = \log 3a + \log 2a$$

$$2 \log x = \log 6a^2$$

$$\log x = \frac{\log 6}{2} + \frac{\log a^2}{2}$$

$$\log x = \frac{1}{2} \log 6 + \frac{2 \log a}{2}$$

$$\log x = \frac{1}{2} \log 6 + \log a$$

PTS: 2 REF: 011224a2 STA: A2.A.19 TOP: Properties of Logarithms


KEY: splitting logs

25 ANS: 3

$$\frac{3y}{2y-6} + \frac{9}{6-2y} = \frac{3y}{2y-6} - \frac{9}{2y-6} = \frac{3y-9}{2y-6} = \frac{3(y-3)}{2(y-3)} = \frac{3}{2}$$

PTS: 2 REF: 011325a2 STA: A2.A.16 TOP: Addition and Subtraction of Rationals

- 26 ANS: 1 PTS: 2 REF: 061420a2 STA: A2.A.34
TOP: Sigma Notation
- 27 ANS: 4 PTS: 2 REF: 011406a2 STA: A2.S.1
TOP: Analysis of Data
- 28 ANS: 4 PTS: 2 REF: 061402a2 STA: A2.A.8
TOP: Negative and Fractional Exponents
- 29 ANS: 3



- PTS: 2 REF: 011207a2 STA: A2.A.71 TOP: Graphing Trigonometric Functions
- 30 ANS: 3

$$-\sqrt{2} \sec x = 2$$

$$\sec x = -\frac{2}{\sqrt{2}}$$

$$\cos x = -\frac{\sqrt{2}}{2}$$

$$x = 135, 225$$

- PTS: 2 REF: 011322a2 STA: A2.A.68 TOP: Trigonometric Equations
KEY: reciprocal functions
- 31 ANS: 3

$$h(-8) = \frac{1}{2}(-8) - 2 = -4 - 2 = -6. \quad g(-6) = \frac{1}{2}(-6) + 8 = -3 + 8 = 5$$

- PTS: 2 REF: 011403a2 STA: A2.A.42 TOP: Compositions of Functions
KEY: numbers
- 32 ANS: 3

$$\log 4m^2 = \log 4 + \log m^2 = \log 4 + 2 \log m$$

- PTS: 2 REF: 061321a2 STA: A2.A.19 TOP: Properties of Logarithms
KEY: splitting logs
- 33 ANS: 3 PTS: 2 REF: 061224a2 STA: A2.A.63
TOP: Domain and Range
- 34 ANS: 1 PTS: 2 REF: 011314a2 STA: A2.N.3
TOP: Operations with Polynomials

35 ANS: 4

$$\frac{2\pi}{b} = 30$$

$$b = \frac{\pi}{15}$$

PTS: 2 REF: 011227a2 STA: A2.A.72

TOP: Identifying the Equation of a Trigonometric Graph

36 ANS: 2

$$\frac{1 - \frac{4}{x}}{1 - \frac{2}{x} - \frac{8}{x^2}} \times \frac{x^2}{x^2} = \frac{x^2 - 4x}{x^2 - 2x - 8} = \frac{x(x-4)}{(x-4)(x+2)} = \frac{x}{x+2}$$

PTS: 2 REF: 061305a2 STA: A2.A.17 TOP: Complex Fractions

37 ANS: 2

$$\sqrt{2x-4} = x-2$$

$$2x-4 = x^2 - 4x + 4$$

$$0 = x^2 - 6x + 8$$

$$0 = (x-4)(x-2)$$

$$x = 4, 2$$

PTS: 2 REF: 061406a2 STA: A2.A.22 TOP: Solving Radicals

KEY: extraneous solutions

38 ANS: 1 PTS: 2 REF: 011310a2 STA: A2.S.9

TOP: Differentiating Permutations and Combinations

39 ANS: 4

$$g\left(\frac{1}{2}\right) = \frac{1}{\frac{1}{2}} = 2. \quad f(2) = 4(2) - 2^2 = 4$$

PTS: 2 REF: 011204a2 STA: A2.A.42 TOP: Compositions of Functions

KEY: numbers

40 ANS: 2 PTS: 2 REF: 061218a2 STA: A2.A.43

TOP: Defining Functions

41 ANS: 4 PTS: 2 REF: 061318a2 STA: A2.A.49

TOP: Equations of Circles

42 ANS: 3 PTS: 2 REF: 061219a2 STA: A2.N.8

TOP: Conjugates of Complex Numbers

43 ANS: 2 PTS: 2 REF: 011417a2 STA: A2.S.9

TOP: Differentiating Permutations and Combinations

44 ANS: 1 PTS: 2 REF: 061409a2 STA: A2.A.38

TOP: Defining Functions KEY: graphs

45 ANS: 1

$$2x - 1 > 5. \quad 2x - 1 < -5$$

$$2x > 6 \quad 2x > -4$$

$$x > 3 \quad x < -2$$

PTS: 2

REF: 061307a2

STA: A2.A.1

TOP: Absolute Value Inequalities

KEY: graph

46 ANS: 3

$$(3i)(2i)^2(m+i)$$

$$(3i)(4i^2)(m+i)$$

$$(3i)(-4)(m+i)$$

$$(-12i)(m+i)$$

$$-12mi - 12i^2$$

$$-12mi + 12$$

PTS: 2

REF: 061319a2

STA: A2.N.9

TOP: Multiplication and Division of Complex Numbers

47 ANS: 1

(4) shows the strongest linear relationship, but if $r < 0$, $b < 0$. The Regents announced that a correct solution was not provided for this question and all students should be awarded credit.

PTS: 2

REF: 011223a2

STA: A2.S.8

TOP: Correlation Coefficient

48 ANS: 1

$$\text{If } \sin x = 0.8, \text{ then } \cos x = 0.6. \quad \tan \frac{1}{2}x = \sqrt{\frac{1-0.6}{1+0.6}} = \sqrt{\frac{0.4}{1.6}} = 0.5.$$

PTS: 2

REF: 061220a2

STA: A2.A.77

TOP: Half Angle Identities

49 ANS: 3

$$\sqrt[3]{4^3 a^{15}} = 4a^5 \sqrt[3]{a}$$

PTS: 2

REF: 061204a2

STA: A2.A.13

TOP: Simplifying Radicals

KEY: index > 2

50 ANS: 1

PTS: 2

REF: 061401a2

STA: A2.S.2

TOP: Analysis of Data

51 ANS: 1

$$20(-2) = x(-2x + 2)$$

$$-40 = -2x^2 + 2x$$

$$2x^2 - 2x - 40 = 0$$

$$x^2 - x - 20 = 0$$

$$(x + 4)(x - 5) = 0$$

$$x = -4, 5$$

PTS: 2

REF: 011321a2

STA: A2.A.5

TOP: Inverse Variation

52 ANS: 1

PTS: 2

REF: 011313a2

STA: A2.A.39

TOP: Domain and Range

KEY: real domain

53 ANS: 3

$$34.1\% + 19.1\% = 53.2\%$$

PTS: 2

REF: 011212a2

STA: A2.S.5

TOP: Normal Distributions

KEY: probability

54 ANS: 1

$$2 \cdot \frac{180}{\pi} = \frac{360}{\pi}$$

PTS: 2

REF: 011220a2

STA: A2.M.2

TOP: Radian Measure

KEY: degrees

55 ANS: 3

PTS: 2

REF: 061418a2

STA: A2.A.51

TOP: Domain and Range

56 ANS: 1

$$5 \cdot \frac{180}{\pi} \approx 286$$

PTS: 2

REF: 011427a2

STA: A2.M.2

TOP: Radian Measure

KEY: degrees

57 ANS: 4

$$\frac{x}{x - \sqrt{x}} \times \frac{x + \sqrt{x}}{x + \sqrt{x}} = \frac{x^2 + x\sqrt{x}}{x^2 - x} = \frac{x(x + \sqrt{x})}{x(x - 1)} = \frac{x + \sqrt{x}}{x - 1}$$

PTS: 2

REF: 061325a2

STA: A2.A.15

TOP: Rationalizing Denominators

KEY: index = 2

58 ANS: 3

$$2! \cdot 2! \cdot 2! = 8$$

PTS: 2

REF: 061425a2

STA: A2.S.10

TOP: Permutations

59 ANS: 2

PTS: 2

REF: 061322a2

STA: A2.A.73

TOP: Law of Sines

KEY: modeling

60 ANS: 2

$$\frac{5}{\sin 32} = \frac{8}{\sin E} \quad 57.98 + 32 < 180$$

$$E \approx 57.98 \quad (180 - 57.98) + 32 < 180$$

PTS: 2 REF: 011419a2 STA: A2.A.75 TOP: Law of Sines - The Ambiguous Case

61 ANS: 4 PTS: 2 REF: 061207a2 STA: A2.A.19

TOP: Properties of Logarithms KEY: antilogarithms

62 ANS: 1

$$\sin(180 + x) = (\sin 180)(\cos x) + (\cos 180)(\sin x) = 0 + (-\sin x) = -\sin x$$

PTS: 2 REF: 011318a2 STA: A2.A.76 TOP: Angle Sum and Difference Identities

KEY: identities

63 ANS: 3

$$42 = \frac{1}{2}(a)(8)\sin 61$$

$$42 \approx 3.5a$$

$$12 \approx a$$

PTS: 2 REF: 011316a2 STA: A2.A.74 TOP: Using Trigonometry to Find Area

KEY: basic

64 ANS: 3 PTS: 2 REF: 061407a2 STA: A2.N.3

TOP: Operations with Polynomials

65 ANS: 4

$${}_3C_2 \left(\frac{5}{8} \right)^2 \left(\frac{3}{8} \right)^1 = \frac{225}{512}$$

PTS: 2 REF: 011221a2 STA: A2.S.15 TOP: Binomial Probability

KEY: spinner

66 ANS: 2 PTS: 2 REF: 011208a2 STA: A2.A.67

TOP: Proving Trigonometric Identities

67 ANS: 1

$$(4a + 4) - (2a + 1) = 2a + 3$$

PTS: 2 REF: 011401a2 STA: A2.A.30 TOP: Sequences

68 ANS: 4 PTS: 2 REF: 061303a2 STA: A2.A.43

TOP: Defining Functions

69 ANS: 3

$${}_8C_3 \cdot x^{8-3} \cdot (-2)^3 = 56x^5 \cdot (-8) = -448x^5$$

PTS: 2 REF: 011308a2 STA: A2.A.36 TOP: Binomial Expansions


70 ANS: 3

$$\frac{4}{-2} = -2$$

PTS: 2 REF: 011304a2 STA: A2.A.31 TOP: Sequences

- 71 ANS: 4 PTS: 1 REF: 011312a2 STA: A2.A.56
TOP: Determining Trigonometric Functions KEY: degrees, common angles
- 72 ANS: 3
 $\sqrt{9} \sqrt{-1} \sqrt{2} - \sqrt{16} \sqrt{-1} \sqrt{2} = 3i\sqrt{2} - 4i\sqrt{2} = -i\sqrt{2}$
- PTS: 2 REF: 061404a2 STA: A2.N.6 TOP: Square Roots of Negative Numbers
- 73 ANS: 4 PTS: 2 REF: 011219a2 STA: A2.A.52
TOP: Properties of Graphs of Functions and Relations
- 74 ANS: 1 PTS: 2 REF: 061408a2 STA: A2.A.24
TOP: Completing the Square
- 75 ANS: 1
$$\frac{6}{\sin 35} = \frac{10}{\sin N}$$
$$N \approx 73$$
$$73 + 35 < 180$$
$$(180 - 73) + 35 < 180$$
- PTS: 2 REF: 061226a2 STA: A2.A.75 TOP: Law of Sines - The Ambiguous Case
- 76 ANS: 3
$$x + y = 5 \quad . \quad -5 + y = 5$$
$$y = -x + 5 \quad y = 10$$
$$(x + 3)^2 + (-x + 5 - 3)^2 = 53$$
$$x^2 + 6x + 9 + x^2 - 4x + 4 = 53$$
$$2x^2 + 2x - 40 = 0$$
$$x^2 + x - 20 = 0$$
$$(x + 5)(x - 4) = 0$$
$$x = -5, 4$$
- PTS: 2 REF: 011302a2 STA: A2.A.3 TOP: Quadratic-Linear Systems
KEY: equations
- 77 ANS: 1 PTS: 2 REF: 061316a2 STA: A2.S.8
TOP: Correlation Coefficient
- 78 ANS: 4
$$2x^2 - 7x - 5 = 0$$
$$\frac{c}{a} = \frac{-5}{2}$$
- PTS: 2 REF: 061414a2 STA: A2.A.20 TOP: Roots of Quadratics

79 ANS: 2



$$\sin^{-1}\left(\frac{8}{17}\right) \text{ DMS}$$

$$28^{\circ}4'20.953''$$

$$\sin S = \frac{8}{17}$$

$$S = \sin^{-1} \frac{8}{17}$$

$$S \approx 28^{\circ}4'$$

PTS: 2 REF: 061311a2 STA: A2.A.55 TOP: Trigonometric Ratios

80 ANS: 3

$$x^2 + y^2 - 16x + 6y + 53 = 0$$

$$x^2 - 16x + 64 + y^2 + 6y + 9 = -53 + 64 + 9$$

$$(x - 8)^2 + (y + 3)^2 = 20$$

PTS: 2 REF: 011415a2 STA: A2.A.47 TOP: Equations of Circles

81 ANS: 3

sum of the roots, $\frac{-b}{a} = \frac{-(-9)}{4} = \frac{9}{4}$. product of the roots, $\frac{c}{a} = \frac{3}{4}$

PTS: 2 REF: 061208a2 STA: A2.A.21 TOP: Roots of Quadratics

KEY: basic

82 ANS: 4

$$4 + 3(2 - x) + 3(3 - x) + 3(4 - x) + 3(5 - x)$$

$$4 + 6 - 3x + 9 - 3x + 12 - 3x + 15 - 3x$$

$$46 - 12x$$

PTS: 2 REF: 061315a2 STA: A2.N.10 TOP: Sigma Notation

KEY: advanced

83 ANS: 4

$$\frac{4 \cdot 0 + 6 \cdot 1 + 10 \cdot 2 + 0 \cdot 3 + 4k + 2 \cdot 5}{4 + 6 + 10 + 0 + k + 2} = 2$$

$$\frac{4k + 36}{k + 22} = 2$$

$$4k + 36 = 2k + 44$$

$$2k = 8$$

$$k = 4$$

PTS: 2 REF: 061221a2 STA: A2.S.3 TOP: Average Known with Missing Data

84 ANS: 2

$$\text{sum: } \frac{-b}{a} = \frac{4}{6} = \frac{2}{3}, \quad \text{product: } \frac{c}{a} = \frac{-12}{6} = -2$$

PTS: 2 REF: 011209a2 STA: A2.A.20 TOP: Roots of Quadratics

85 ANS: 3

$$3x^3 - 5x^2 - 48x + 80$$

$$x^2(3x - 5) - 16(3x - 5)$$

$$(x^2 - 16)(3x - 5)$$

$$(x + 4)(x - 4)(3x - 5)$$

PTS: 2 REF: 011317a2 STA: A2.A.7 TOP: Factoring by Grouping

86 ANS: 4

$$\log 2x^3 = \log 2 + \log x^3 = \log 2 + 3 \log x$$

PTS: 2 REF: 061426a2 STA: A2.A.19 TOP: Properties of Logarithms

KEY: splitting logs

87 ANS: 1

$$\frac{1}{7 - \sqrt{11}} \cdot \frac{7 + \sqrt{11}}{7 + \sqrt{11}} = \frac{7 + \sqrt{11}}{49 - 11} = \frac{7 + \sqrt{11}}{38}$$

PTS: 2 REF: 011404a2 STA: A2.N.5 TOP: Rationalizing Denominators

88 ANS: 1

$$\frac{{}_6P_6}{3!2!} = \frac{720}{12} = 60$$

PTS: 2 REF: 011324a2 STA: A2.S.10 TOP: Permutations

89 ANS: 1

PTS: 2 REF: 011416a2 STA: A2.A.39

TOP: Domain and Range

KEY: real domain

90 ANS: 3

PTS: 2 REF: 061412a2 STA: A2.A.60

TOP: Finding the Terminal Side of an Angle

91 ANS: 1

PTS: 2 REF: 011306a2 STA: A2.A.8

TOP: Negative and Fractional Exponents


92 ANS: 1

$$\frac{9}{\sin A} = \frac{10}{\sin 70^\circ}. \quad 58^\circ + 70^\circ \text{ is possible. } 122^\circ + 70^\circ \text{ is not possible.}$$

$$A = 58$$

PTS: 2 REF: 011210a2 STA: A2.A.75 TOP: Law of Sines - The Ambiguous Case

93 ANS: 4



PTS: 2 REF: 061217a2 STA: A2.A.66 TOP: Determining Trigonometric Functions

94 ANS: 2 PTS: 2 REF: 011225a2 STA: A2.A.43

TOP: Defining Functions

95 ANS: 2

$$\frac{30}{(x+3)(x-3)} + \frac{(x+3)(x-3)}{(x+3)(x-3)} = \frac{5(x+3)}{(x-3)(x+3)} \quad 3 \text{ is an extraneous root.}$$

$$30 + x^2 - 9 = 5x + 15$$

$$x^2 - 5x + 6 = 0$$

$$(x-3)(x-2) = 0$$

$$x = 2$$

PTS: 2 REF: 061417a2 STA: A2.A.23 TOP: Solving Rationals

KEY: rational solutions

96 ANS: 1

$$f(g(x)) = 2(x+5)^2 - 3(x+5) + 1 = 2(x^2 + 10x + 25) - 3x - 15 + 1 = 2x^2 + 17x + 36$$

PTS: 2 REF: 061419a2 STA: A2.A.42 TOP: Compositions of Functions

KEY: variables

97 ANS: 3

$$\sqrt[3]{6a^4b^2} + \sqrt[3]{(27 \cdot 6)a^4b^2}$$

$$a^3\sqrt[3]{6ab^2} + 3a^3\sqrt[3]{6ab^2}$$

$$4a^3\sqrt[3]{6ab^2}$$

PTS: 2 REF: 011319a2 STA: A2.N.2 TOP: Operations with Radicals

98 ANS: 4

$$\frac{x^2 + 9x - 22}{x^2 - 121} \div (2 - x) = \frac{(x+11)(x-2)}{(x+11)(x-11)} \cdot \frac{-1}{x-2} = \frac{-1}{x-11}$$

PTS: 2 REF: 011423a2 STA: A2.A.16 TOP: Multiplication and Division of Rationals

KEY: Division

99 ANS: 3

$$\frac{4x-5}{3} > 1 \text{ or } \frac{4x-5}{3} < -1$$

$$4x-5 > 3 \quad 4x-5 < -3$$

$$4x > 8 \quad 4x < 2$$

$$x > 2 \quad x < \frac{1}{2}$$

PTS: 2 REF: 061209a2 STA: A2.A.1 TOP: Absolute Value Inequalities

KEY: graph

100 ANS: 4 PTS: 2 REF: 061427a2 STA: A2.A.63

TOP: Domain and Range

101 ANS: 2 PTS: 2 REF: 061205a2 STA: A2.A.34

TOP: Sigma Notation

102 ANS: 3

$$S_8 = \frac{3(1 - (-4)^8)}{1 - (-4)} = \frac{196,605}{5} = -39,321$$

PTS: 2 REF: 061304a2 STA: A2.A.35 TOP: Summations

KEY: geometric

103 ANS: 4

$$x^2(x+2) - (x+2)$$

$$(x^2 - 1)(x+2)$$

$$(x+1)(x-1)(x+2)$$

PTS: 2 REF: 011426a2 STA: A2.A.7 TOP: Factoring by Grouping

104 ANS: 3 PTS: 2 REF: 061308ge STA: A2.A.51

TOP: Domain and Range

105 ANS: 1 PTS: 2 REF: 061210a2 STA: A2.A.9

TOP: Negative Exponents

106 ANS: 2

$$2^2 \cdot 3 = 12 \quad 6^2 d = 12$$

$$4^2 \cdot \frac{3}{4} = 12 \quad 36d = 12$$

$$d = \frac{1}{3}$$

PTS: 2 REF: 061310a2 STA: A2.A.5 TOP: Inverse Variation

107 ANS: 4 PTS: 2 REF: 061411a2 STA: A2.A.30

TOP: Sequences

108 ANS: 1

$$10 \cdot \frac{3}{2} = \frac{3}{5}p$$

$$15 = \frac{3}{5}p$$

$$25 = p$$

PTS: 2

REF: 011226a2

STA: A2.A.5

TOP: Inverse Variation

109 ANS: 2

$$\text{If } \sin A = -\frac{7}{25}, \cos A = \frac{24}{25}, \text{ and } \tan A = \frac{\sin A}{\cos A} = \frac{-\frac{7}{25}}{\frac{24}{25}} = -\frac{7}{24}$$

PTS: 2

REF: 011413a2

STA: A2.A.64

TOP: Using Inverse Trigonometric Functions

KEY: advanced

110 ANS: 2

$$320 = 10(2)^{\frac{t}{60}}$$

$$32 = (2)^{\frac{t}{60}}$$

$$\log 32 = \log(2)^{\frac{t}{60}}$$

$$\log 32 = \frac{t \log 2}{60}$$

$$\frac{60 \log 32}{\log 2} = t$$

$$300 = t$$

PTS: 2

REF: 011205a2

STA: A2.A.6

TOP: Exponential Growth

111 ANS: 3

$$S_n = \frac{n}{2} [2a + (n-1)d] = \frac{19}{2} [2(3) + (19-1)7] = 1254$$

PTS: 2

REF: 011202a2

STA: A2.A.35

TOP: Summations

KEY: arithmetic

112 ANS: 2

$$\bar{x} \pm \sigma$$

$$153 \pm 22$$

$$131 - 175$$

PTS: 2

REF: 011307a2

STA: A2.S.5

TOP: Normal Distributions

KEY: interval

113 ANS: 4

$$(a-1)^2 + (a-2)^2 + (a-3)^2 + (a-4)^2$$

$$(a^2 - 2a + 1) + (a^2 - 4a + 4) + (a^2 - 6a + 9) + (a^2 - 8a + 16)$$

$$4a^2 - 20a + 30$$

PTS: 2 REF: 011414a2 STA: A2.N.10 TOP: Sigma Notation
KEY: advanced

114 ANS: 2

$$\frac{1}{2}(22)(13)\sin 55 \approx 117$$

PTS: 2 REF: 061403a2 STA: A2.A.74 TOP: Using Trigonometry to Find Area
KEY: basic

115 ANS: 1 PTS: 2 REF: 011320a2 STA: A2.A.72

TOP: Identifying the Equation of a Trigonometric Graph

116 ANS: 3 PTS: 2 REF: 061416a2 STA: A2.A.12

TOP: Evaluating Exponential Expressions

117 ANS: 3

$$\frac{-b}{a} = \frac{-(-4)}{1} = 4. \text{ If the sum is 4, the roots must be 7 and } -3.$$

PTS: 2 REF: 011418a2 STA: A2.A.21 TOP: Roots of Quadratics
KEY: advanced

118 ANS: 1 PTS: 2 REF: 061324a2 STA: A2.A.9

TOP: Negative Exponents

119 ANS: 2

The binomials are conjugates, so use FL.

PTS: 2 REF: 011206a2 STA: A2.N.3 TOP: Operations with Polynomials

120 ANS: 1 PTS: 2 REF: 061317a2 STA: A2.S.9

TOP: Differentiating Permutations and Combinations

121 ANS: 2 PTS: 2 REF: 011315a2 STA: A2.A.55

TOP: Trigonometric Ratios

122 ANS: 3

As originally written, alternatives (2) and (3) had no domain restriction, so that both were correct.


PTS: 2 REF: 061405a2 STA: A2.A.52
TOP: Properties of Graphs of Functions and Relations

123 ANS: 1

The binomials are conjugates, so use FL.

PTS: 2 REF: 061201a2 STA: A2.N.3 TOP: Operations with Polynomials

124 ANS: 4



$$2 \cos \theta = 1$$

$$\cos \theta = \frac{1}{2}$$

$$\theta = \cos^{-1} \frac{1}{2} = 60, 300$$

PTS: 2 REF: 061203a2 STA: A2.A.68 TOP: Trigonometric Equations
KEY: basic

125 ANS: 3 PTS: 2 REF: 011305a2 STA: A2.A.38
TOP: Defining Functions KEY: graphs

126 ANS: 1 PTS: 2 REF: 061223a2 STA: A2.S.15
TOP: Binomial Probability KEY: modeling

127 ANS: 2
 $\log 9 - \log 20$

$$\log 3^2 - \log(10 \cdot 2)$$

$$2 \log 3 - (\log 10 + \log 2)$$

$$2b - (1 + a)$$

$$2b - a - 1$$

PTS: 2 REF: 011326a2 STA: A2.A.19 TOP: Properties of Logarithms
KEY: expressing logs algebraically


128 ANS: 3
 ${}_3C_2(2x^4)^1(-y)^2 = 6x^4y^2$

PTS: 2 REF: 011215a2 STA: A2.A.36 TOP: Binomial Expansions

129 ANS: 4
 $(x+i)^2 - (x-i)^2 = x^2 + 2xi + i^2 - (x^2 - 2xi + i^2) = 4xi$

PTS: 2 REF: 011327a2 STA: A2.N.9
TOP: Multiplication and Division of Complex Numbers

130 ANS: 1



PTS: 2 REF: 061225a2 STA: A2.S.8 TOP: Correlation Coefficient

131 ANS: 2

$$\frac{2\pi}{6} = \frac{\pi}{3}$$

PTS: 2 REF: 061413a2 STA: A2.A.69

TOP: Properties of Graphs of Trigonometric Functions

KEY: period

132 ANS: 2

$$\cos(x - y) = \cos x \cos y + \sin x \sin y$$

$$= b \cdot b + a \cdot a$$

$$= b^2 + a^2$$

PTS: 2 REF: 061421a2 STA: A2.A.76

TOP: Angle Sum and Difference Identities

KEY: simplifying

133 ANS: 2

$$\frac{x^{-1} + 1}{x + 1} = \frac{\frac{1}{x} + 1}{x + 1} = \frac{\frac{1 + x}{x}}{x + 1} = \frac{1}{x}$$

PTS: 2 REF: 011211a2 STA: A2.A.9

TOP: Negative Exponents

134 ANS: 2

$$\tan 30 = \frac{\sqrt{3}}{3} \cdot \text{Arc cos } \frac{\sqrt{3}}{k} = 30$$

$$\frac{\sqrt{3}}{k} = \cos 30$$

$$k = 2$$

PTS: 2 REF: 061323a2 STA: A2.A.64

TOP: Using Inverse Trigonometric Functions

KEY: advanced

135 ANS: 1

PTS: 2

REF: 011402a2

STA: A2.A.8

TOP: Negative and Fractional Exponents

136 ANS: 4

PTS: 2

REF: 011201a2

STA: A2.S.2

TOP: Analysis of Data

137 ANS: 3

$$3x^5 - 48x = 0$$

$$3x(x^4 - 16) = 0$$

$$3x(x^2 + 4)(x^2 - 4) = 0$$

$$3x(x^2 + 4)(x + 2)(x - 2) = 0$$

PTS: 2 REF: 011216a2 STA: A2.A.26

TOP: Solving Polynomial Equations

138 ANS: 4

$${}_{15}C_5 = 3,003. \quad {}_{25}C_5 = {}_{25}C_{20} = 53,130. \quad {}_{25}C_{15} = 3,268,760.$$

PTS: 2 REF: 061227a2 STA: A2.S.11

TOP: Combinations

139 ANS: 4

$$8^{3k+4} = 4^{2k-1}$$

$$(2^3)^{3k+4} = (2^2)^{2k-1}$$

$$2^{9k+12} = 2^{4k-2}$$

$$9k+12 = 4k-2$$

$$5k = -14$$

$$k = -\frac{14}{5}$$

PTS: 2 REF: 011309a2 STA: A2.A.27 TOP: Exponential Equations
KEY: common base not shown

140 ANS: 3 PTS: 2 REF: 011422a2 STA: A2.A.54
TOP: Graphing Logarithmic Functions

141 ANS: 3

$$5000 \left(1 + \frac{.03}{4} \right)^{4 \cdot 5} = 5000(1.0075)^{20} \approx 5805.92$$

PTS: 2 REF: 011410a2 STA: A2.A.12 TOP: Evaluating Exponential Expressions
142 ANS: 2 PTS: 2 REF: 061216a2 STA: A2.A.42
TOP: Compositions of Functions KEY: variables

143 ANS: 3

$$\frac{x+16}{x-2} - \frac{7(x-2)}{x-2} \leq 0 \quad -6x+30=0 \quad x-2=0. \text{ Check points such that } x < 2, 2 < x < 5, \text{ and } x > 5. \text{ If } x = 1,$$

$$\frac{-6x+30}{x-2} \leq 0 \quad -6x = -30 \quad x = 2$$

$$\frac{-6(1)+30}{1-2} = \frac{24}{-1} = -24, \text{ which is less than } 0. \text{ If } x = 3, \frac{-6(3)+30}{3-2} = \frac{12}{1} = 12, \text{ which is greater than } 0. \text{ If } x = 6,$$

$$\frac{-6(6)+30}{6-2} = \frac{-6}{4} = -\frac{3}{2}, \text{ which is less than } 0.$$

PTS: 2 REF: 011424a2 STA: A2.A.23 TOP: Rational Inequalities
144 ANS: 2

$$\frac{\cot x}{\csc x} = \frac{\frac{\cos x}{\sin x}}{\frac{1}{\sin x}} = \cos x$$

PTS: 2 REF: 061410a2 STA: A2.A.58 TOP: Reciprocal Trigonometric Relationships
145 ANS: 2
Since the coefficient of t is greater than 0, $r > 0$.

PTS: 2 REF: 011303a2 STA: A2.S.8 TOP: Correlation Coefficient

146 ANS: 4

$$x = 2y. \quad y^2 - (3y)^2 + 32 = 0 \quad . \quad x = 3(-2) = -6$$

$$y^2 - 9y^2 = -32$$

$$-8y^2 = -32$$

$$y^2 = 4$$

$$y = \pm 2$$

PTS: 2

REF: 061312a2

STA: A2.A.3

TOP: Quadratic-Linear Systems

KEY: equations

147 ANS: 3

If $\csc P > 0$, $\sin P > 0$. If $\cot P < 0$ and $\sin P > 0$, $\cos P < 0$

PTS: 2

REF: 061320a2

STA: A2.A.60

TOP: Finding the Terminal Side of an Angle

148 ANS: 4

$$\left(\sqrt[3]{27x^2}\right)\left(\sqrt[3]{16x^4}\right) = \sqrt[3]{3^3 \cdot 2^4 \cdot x^6} = 3 \cdot 2 \cdot x^{2\frac{2}{3}}\sqrt{2} = 6x^{2\frac{2}{3}}\sqrt{2}$$

PTS: 2

REF: 011421a2

STA: A2.N.2

TOP: Operations with Radicals

149 ANS: 1

PTS: 2

REF: 061202a2

STA: A2.A.51

TOP: Domain and Range

150 ANS: 2

PTS: 2

REF: 011407a2

STA: A2.A.43

TOP: Defining Functions

151 ANS: 4

$$\cos 2A = 1 - 2\sin^2 A = 1 - 2\left(\frac{1}{3}\right)^2 = 1 - \frac{2}{9} = \frac{7}{9}$$

PTS: 2


REF: 011311a2

STA: A2.A.77

TOP: Double Angle Identities

KEY: evaluating

152 ANS: 1



A calculator screen showing the expression $\cos\left(\frac{5\pi}{6}\right)$ and the result -1.154700538 .

PTS: 2

REF: 011203a2

STA: A2.A.66

TOP: Determining Trigonometric Functions

153 ANS: 3

$$20 \cdot 2 = -5t$$

$$-8 = t$$

PTS: 2

REF: 011412a2

STA: A2.A.5

TOP: Inverse Variation

154 ANS: 1

$$\frac{2\pi}{b} = 4\pi$$

$$b = \frac{1}{2}$$

PTS: 2 REF: 011425a2 STA: A2.A.69

TOP: Properties of Graphs of Trigonometric Functions

KEY: period

155 ANS: 2

$$\frac{-\frac{3}{32}a^3b^4}{\frac{1}{64}a^5b^3} = -\frac{6b}{a^2}$$

PTS: 2 REF: 061326a2 STA: A2.A.31

TOP: Sequences

156 ANS: 1

$$\cos(A - B) = \left(\frac{5}{13}\right)\left(-\frac{3}{5}\right) + \left(\frac{12}{13}\right)\left(\frac{4}{5}\right) = -\frac{15}{65} + \frac{48}{65} = \frac{33}{65}$$

PTS: 2 REF: 011214a2 STA: A2.A.76

TOP: Angle Sum and Difference Identities

KEY: evaluating

157 ANS: 2

PTS: 2

REF: 011213a2

STA: A2.N.8

TOP: Conjugates of Complex Numbers

158 ANS: 3

$${}_6C_3 \left(\frac{x}{2}\right)^3 (-2y)^3 = 20 \cdot \frac{x^3}{8} \cdot -8y^3 = -20x^3y^3$$

PTS: 2 REF: 061215a2 STA: A2.A.36

TOP: Binomial Expansions

159 ANS: 3

$$1000 = 500e^{.05t}$$

$$2 = e^{.05t}$$

$$\ln 2 = \ln e^{.05t}$$

$$\frac{\ln 2}{.05} = \frac{.05t \cdot \ln e}{.05}$$

$$13.9 \approx t$$

PTS: 2 REF: 061313a2 STA: A2.A.6

TOP: Exponential Growth

160 ANS: 4


$$\frac{13}{\sin 40} = \frac{20}{\sin M} \cdot 81 + 40 < 180. \quad (180 - 81) + 40 < 180$$

$$M \approx 81$$

PTS: 2 REF: 061327a2 STA: A2.A.75

TOP: Law of Sines - The Ambiguous Case

161 ANS: 4



PTS: 2

REF: 061222a2

STA: A2.A.50

TOP: Solving Polynomial Equations

162 ANS: 1

PTS: 2

REF: 061211a2

STA: A2.A.54

TOP: Graphing Logarithmic Functions

Algebra 2/Trigonometry Multiple Choice Regents Exam Questions Answer Section

163 ANS: 2

$$x^2 + 2 = 6x$$

$$x^2 - 6x = -2$$

$$x^2 - 6x + 9 = -2 + 9$$

$$(x - 3)^2 = 7$$

PTS: 2

REF: 011116a2

STA: A2.A.24

TOP: Completing the Square

164 ANS: 4

$$\frac{3 \pm \sqrt{(-3)^2 - 4(1)(-9)}}{2(1)} = \frac{3 \pm \sqrt{45}}{2} = \frac{3 \pm 3\sqrt{5}}{2}$$

PTS: 2

REF: 061009a2

STA: A2.A.25

TOP: Quadratics with Irrational Solutions

165 ANS: 1

PTS: 2

REF: 061013a2

STA: A2.A.38

TOP: Defining Functions

166 ANS: 1

PTS: 2

REF: 061019a2

STA: A2.N.7

TOP: Imaginary Numbers

167 ANS: 4

PTS: 2

REF: 081005a2

STA: A2.A.60

TOP: Unit Circle

168 ANS: 2

$$f(10) = \frac{-10}{(-10)^2 - 16} = \frac{-10}{84} = -\frac{5}{42}$$

PTS: 2

REF: 061102a2

STA: A2.A.41

TOP: Functional Notation

169 ANS: 4

$$s = \theta r = 2 \cdot 4 = 8$$

PTS: 2

REF: fall0922a2

STA: A2.A.61

TOP: Arc Length

KEY: arc length

170 ANS: 3

$$\frac{59.2}{\sin 74} = \frac{60.3}{\sin C} \quad 180 - 78.3 = 101.7$$

$$C \approx 78.3$$

PTS: 2

REF: 081006a2

STA: A2.A.75

TOP: Law of Sines - The Ambiguous Case

171 ANS: 4

$$y - 2 \sin \theta = 3$$

$$y = 2 \sin \theta + 3$$

$$f(\theta) = 2 \sin \theta + 3$$

PTS: 2

REF: fall0927a2

STA: A2.A.40

TOP: Functional Notation

172 ANS: 3

$$x^2 - 3x - 10 > 0 \quad \text{or}$$

$$(x - 5)(x + 2) > 0 \quad x - 5 < 0 \text{ and } x + 2 < 0$$

$$x - 5 > 0 \text{ and } x + 2 > 0 \quad x < 5 \text{ and } x < -2$$

$$x > 5 \text{ and } x > -2 \quad x < -2$$

$$x > 5$$

PTS: 2

REF: 011115a2

STA: A2.A.4

TOP: Quadratic Inequalities

KEY: one variable

173 ANS: 2

$$4^{2x+5} = 8^{3x}$$

$$\left(2^2\right)^{2x+5} = \left(2^3\right)^{3x}$$

$$2^{4x+10} = 2^{9x}$$

$$4x + 10 = 9x$$

$$10 = 5x$$

$$2 = x$$

PTS: 2

REF: 061105a2

STA: A2.A.27

TOP: Exponential Equations

KEY: common base not shown

174 ANS: 1

$$6x - 7 \leq 5 \quad 6x - 7 \geq -5$$

$$6x \leq 12 \quad 6x \geq 2$$

$$x \leq 2 \quad x \geq \frac{1}{3}$$

PTS: 2

REF: fall0905a2

STA: A2.A.1

TOP: Absolute Value Inequalities

KEY: graph

175 ANS: 3

$$\sqrt{-300} = \sqrt{100} \sqrt{-1} \sqrt{3}$$

PTS: 2

REF: 061006a2

STA: A2.N.6

TOP: Square Roots of Negative Numbers

176 ANS: 1

$$a_n = -\sqrt{5}(-\sqrt{2})^{n-1}$$

$$a_{15} = -\sqrt{5}(-\sqrt{2})^{15-1} = -\sqrt{5}(-\sqrt{2})^{14} = -\sqrt{5} \cdot 2^7 = -128\sqrt{5}$$


PTS: 2

REF: 061109a2

STA: A2.A.32

TOP: Sequences

177 ANS: 1



$$\tan \theta - \sqrt{3} = 0$$

$$\tan \theta = \sqrt{3}$$

$$\theta = \tan^{-1} \sqrt{3}$$

$$\theta = 60, 240$$

PTS: 2 REF: fall0903a2 STA: A2.A.68 TOP: Trigonometric Equations
KEY: basic

178 ANS: 3

$$3x + 16 = (x + 2)^2 \quad . \quad -4 \text{ is an extraneous solution.}$$

$$3x + 16 = x^2 + 4x + 4$$

$$0 = x^2 + x - 12$$

$$0 = (x + 4)(x - 3)$$

$$x = -4 \quad x = 3$$

PTS: 2 REF: 061121a2 STA: A2.A.22 TOP: Solving Radicals
KEY: extraneous solutions

179 ANS: 1 PTS: 2 REF: 081022a2 STA: A2.A.46
TOP: Transformations with Functions and Relations

180 ANS: 3 PTS: 2 REF: 011119a2 STA: A2.A.52
TOP: Families of Functions

181 ANS: 1

$$2i^2 + 3i^3 = 2(-1) + 3(-i) = -2 - 3i$$

PTS: 2 REF: 081004a2 STA: A2.N.7 TOP: Imaginary Numbers

182 ANS: 1 PTS: 2 REF: fall0914a2 STA: A2.A.9
TOP: Negative and Fractional Exponents

183 ANS: 1

$8 \times 8 \times 7 \times 1 = 448$. The first digit cannot be 0 or 5. The second digit cannot be 5 or the same as the first digit. The third digit cannot be 5 or the same as the first or second digit.

PTS: 2 REF: 011125a2 STA: A2.S.10 TOP: Permutations

184 ANS: 2

$$\left(\frac{w^{-5}}{w^{-9}} \right)^{\frac{1}{2}} = (w^4)^{\frac{1}{2}} = w^2$$

PTS: 2 REF: 081011a2 STA: A2.A.8 TOP: Negative and Fractional Exponents

185 ANS: 2

The roots are $-1, 2, 3$.

PTS: 2

REF: 081023a2

STA: A2.A.50

TOP: Solving Polynomial Equations

186 ANS: 3

$$2\pi \cdot \frac{5}{12} = \frac{10\pi}{12} = \frac{5\pi}{6}$$

PTS: 2

REF: 061125a2

STA: A2.M.1

TOP: Radian Measure

187 ANS: 2

$$(3 - 7i)(3 - 7i) = 9 - 21i - 21i + 49i^2 = 9 - 42i - 49 = -40 - 42i$$

PTS: 2

REF: fall0901a2

STA: A2.N.9

TOP: Multiplication and Division of Complex Numbers

188 ANS: 2

$$x^2 - x - 6 = 3x - 6$$

$$x^2 - 4x = 0$$

$$x(x - 4) = 0$$

$$x = 0, 4$$

PTS: 2

REF: 081015a2

STA: A2.A.3

TOP: Quadratic-Linear Systems

KEY: equations

189 ANS: 1

$$\sqrt{12^2 - 6^2} = \sqrt{108} = \sqrt{36} \sqrt{3} = 6\sqrt{3}. \cot J = \frac{A}{O} = \frac{6}{6\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{\sqrt{3}}{3}$$

PTS: 2

REF: 011120a2

STA: A2.A.55

TOP: Trigonometric Ratios

190 ANS: 4

PTS: 2

REF: 061112a2

STA: A2.A.39

TOP: Domain and Range

KEY: real domain

191 ANS: 1

```


Plot1 Plot2 Plot3
Y1=1/sin(X)
Y2=
Y3=
Y4=
Y5=
Y6=
Y7=

```

```

WINDOW
Xmin=-3.141592...
Xmax=3.1415926...
Xscl=0
Ymin=-10
Ymax=10
Yscl=1
Xres=1

```



PTS: 2

REF: 011123a2

STA: A2.A.71

TOP: Graphing Trigonometric Functions

192 ANS: 2

PTS: 2

REF: 081024a2

STA: A2.N.8

TOP: Conjugates of Complex Numbers

193 ANS: 4

PTS: 2

REF: 061005a2

STA: A2.A.50

TOP: Solving Polynomial Equations

194 ANS: 4

$$9^{3x+1} = 27^{x+2}$$

$$(3^2)^{3x+1} = (3^3)^{x+2}$$

$$3^{6x+2} = 3^{3x+6}$$

$$6x+2 = 3x+6$$

$$3x = 4$$

$$x = \frac{4}{3}$$

PTS: 2

REF: 081008a2

STA: A2.A.27

TOP: Exponential Equations

KEY: common base not shown

195 ANS: 3

$$4^{x^2+4x} = 2^{-6} \quad 2x^2 + 8x = -6$$

$$(2^2)^{x^2+4x} = 2^{-6} \quad 2x^2 + 8x + 6 = 0$$

$$2^{2x^2+8x} = 2^{-6} \quad x^2 + 4x + 3 = 0$$

$$(x+3)(x+1) = 0$$

$$x = -3 \quad x = -1$$

PTS: 2

REF: 061015a2

STA: A2.A.27

TOP: Exponential Equations

KEY: common base shown

196 ANS: 3

PTS: 2

REF: 061114a2

STA: A2.A.38

TOP: Defining Functions

KEY: graphs

197 ANS: 2

PTS: 2

REF: fall0926a2

STA: A2.A.46

TOP: Transformations with Functions and Relations

198 ANS: 2


PTS: 2

REF: 081010a2

STA: A2.A.55

TOP: Trigonometric Ratios

199 ANS: 1



PTS: 2

REF: fall0915a2

STA: A2.S.5

TOP: Normal Distributions

KEY: interval

200 ANS: 4

$$\frac{2\pi}{b} = \frac{2\pi}{\frac{1}{3}} = 6\pi$$

PTS: 2 REF: 061027a2 STA: A2.A.69

TOP: Properties of Graphs of Trigonometric Functions

KEY: period

201 ANS: 4

$$x^{-\frac{2}{5}} = \frac{1}{x^{\frac{2}{5}}} = \frac{1}{\sqrt[5]{x^2}}$$

PTS: 2 REF: 011118a2 STA: A2.A.10

TOP: Fractional Exponents as Radicals

202 ANS: 3

$$\frac{-7 \pm \sqrt{7^2 - 4(2)(-3)}}{2(2)} = \frac{-7 \pm \sqrt{73}}{4}$$

PTS: 2 REF: 081009a2 STA: A2.A.25

TOP: Quadratics with Irrational Solutions

203 ANS: 3

$$\left(\frac{2}{3}\right)^2 + \cos^2 A = 1$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos^2 A = \frac{5}{9} = 2\left(\frac{2}{3}\right)\left(\frac{\sqrt{5}}{3}\right)$$

$$\cos A = +\frac{\sqrt{5}}{3}, \sin A \text{ is acute.} = \frac{4\sqrt{5}}{9}$$

PTS: 2 REF: 011107a2 STA: A2.A.77

TOP: Double Angle Identities

KEY: evaluating

204 ANS: 3

PTS: 2

REF: 061119a2

STA: A2.A.65

TOP: Graphing Trigonometric Functions

205 ANS: 4

PTS: 2

REF: 011111a2

STA: A2.N.8

TOP: Conjugates of Complex Numbers

206 ANS: 3

$$\text{period} = \frac{2\pi}{b} = \frac{2\pi}{3\pi} = \frac{2}{3}$$

PTS: 2 REF: 081026a2 STA: A2.A.70

TOP: Graphing Trigonometric Functions

KEY: recognize

207 ANS: 3

PTS: 2

REF: 061007a2

STA: A2.S.9

TOP: Differentiating Permutations and Combinations

208 ANS: 1

common difference is 2. $b_n = x + 2n$

$$10 = x + 2(1)$$

$$8 = x$$

PTS: 2

REF: 081014a2

STA: A2.A.29

TOP: Sequences

209 ANS: 2

$$6(x^2 - 5) = 6x^2 - 30$$

PTS: 2

REF: 011109a2

STA: A2.A.42

TOP: Compositions of Functions

KEY: variables

210 ANS: 1

PTS: 2

REF: 061004a2

STA: A2.A.52

TOP: Identifying the Equation of a Graph

211 ANS: 3

PTS: 2

REF: 011110a2

STA: A2.A.30

TOP: Sequences

212 ANS: 4

Students entering the library are more likely to spend more time studying, creating bias.


PTS: 2

REF: fall0904a2

STA: A2.S.2

TOP: Analysis of Data

213 ANS: 1



$$\cos K = \frac{5}{6}$$

$$K = \cos^{-1} \frac{5}{6}$$

$$K \approx 33^\circ 33'$$

PTS: 2

REF: 061023a2

STA: A2.A.55

TOP: Trigonometric Ratios

214 ANS: 3

$$\frac{3}{\sqrt{3a^2b}} = \frac{3}{a\sqrt{3b}} \cdot \frac{\sqrt{3b}}{\sqrt{3b}} = \frac{3\sqrt{3b}}{3ab} = \frac{\sqrt{3b}}{ab}$$

PTS: 2

REF: 081019a2

STA: A2.A.15

TOP: Rationalizing Denominators

KEY: index = 2

215 ANS: 3

$$a_n = 5(-2)^{n-1}$$

$$a_{15} = 5(-2)^{15-1} = 81,920$$

PTS: 2

REF: 011105a2

STA: A2.A.32

TOP: Sequences

216 ANS: 4 PTS: 2 REF: 011127a2 STA: A2.S.1
TOP: Analysis of Data

217 ANS: 3

1-Var Stats L1, L2	σx^2 67.31102041
--------------------	-----------------------------

PTS: 2 REF: fall0924a2 STA: A2.S.4 TOP: Dispersion
KEY: range, quartiles, interquartile range, variance

218 ANS: 3

$$\frac{\sin^2 \theta + \cos^2 \theta}{1 - \sin^2 \theta} = \frac{1}{\cos^2 \theta} = \sec^2 \theta$$

PTS: 2 REF: 061123a2 STA: A2.A.58 TOP: Reciprocal Trigonometric Relationships

219 ANS: 1

n	3	4	5	Σ
$-r^2 + r$	$-3^2 + 3 = -6$	$-4^2 + 4 = -12$	$-5^2 + 5 = -20$	-38

PTS: 2 REF: 061118a2 STA: A2.N.10 TOP: Sigma Notation
KEY: basic

220 ANS: 3

$$K = (10)(18)\sin 46 \approx 129$$

PTS: 2 REF: 081021a2 STA: A2.A.74 TOP: Using Trigonometry to Find Area
KEY: parallelograms

221 ANS: 2 PTS: 2 REF: 061108a2 STA: A2.A.52

TOP: Identifying the Equation of a Graph

222 ANS: 2

$$\frac{\frac{x}{4} - \frac{1}{x}}{\frac{1}{2x} + \frac{1}{4}} = \frac{\frac{x^2 - 4}{4x}}{\frac{2x + 4}{8x}} = \frac{(x + 2)(x - 2)}{4x} \times \frac{8x}{2(x + 2)} = x - 2$$

PTS: 2 REF: fall0920a2 STA: A2.A.17 TOP: Complex Fractions

223 ANS: 3

$$75000 = 25000e^{.0475t}$$

$$3 = e^{.0475t}$$

$$\ln 3 = \ln e^{.0475t}$$

$$\frac{\ln 3}{.0475} = \frac{.0475t \cdot \ln e}{.0475}$$

$$23.1 \approx t$$

PTS: 2 REF: 061117a2 STA: A2.A.6 TOP: Exponential Growth

224 ANS: 1 PTS: 2 REF: 061018a2 STA: A2.A.22

TOP: Solving Radicals

KEY: extraneous solutions

225 ANS: 3

$$b^2 - 4ac = (-10)^2 - 4(1)(25) = 100 - 100 = 0$$

PTS: 2 REF: 011102a2 STA: A2.A.2 TOP: Using the Discriminant

KEY: determine nature of roots given equation

226 ANS: 1

$$y \geq x^2 - x - 6$$

$$y \geq (x - 3)(x + 2)$$

PTS: 2 REF: 061017a2 STA: A2.A.4 TOP: Quadratic Inequalities

KEY: two variables

227 ANS: 3

$$\frac{4}{5 - \sqrt{13}} \cdot \frac{5 + \sqrt{13}}{5 + \sqrt{13}} = \frac{4(5 + \sqrt{13})}{25 - 13} = \frac{5 + \sqrt{13}}{3}$$

PTS: 2 REF: 061116a2 STA: A2.N.5 TOP: Rationalizing Denominators

228 ANS: 2

$$8^2 = 64$$

PTS: 2 REF: fall0909a2 STA: A2.A.18 TOP: Evaluating Logarithmic Expressions

229 ANS: 2 PTS: 2 REF: 061011a2 STA: A2.A.10

TOP: Fractional Exponents as Radicals

230 ANS: 2 PTS: 2 REF: 011114a2 STA: A2.N.3

TOP: Operations with Polynomials

231 ANS: 4 PTS: 2 REF: fall0908a2 STA: A2.A.38

TOP: Defining Functions KEY: graphs

232 ANS: 2 PTS: 2 REF: 081003a2 STA: A2.A.51

TOP: Domain and Range

233 ANS: 4

$$12x^4 + 10x^3 - 12x^2 = 2x^2(6x^2 + 5x - 6) = 2x^2(2x + 3)(3x - 2)$$

PTS: 2 REF: 061008a2 STA: A2.A.7 TOP: Factoring Polynomials

KEY: single variable

234 ANS: 4

$$S_n = \frac{n}{2} [2a + (n-1)d] = \frac{21}{2} [2(18) + (21-1)2] = 798$$

PTS: 2 REF: 061103a2 STA: A2.A.35 TOP: Series
KEY: arithmetic

235 ANS: 2

$$f^{-1}(x) = \log_4 x$$

PTS: 2 REF: fall0916a2 STA: A2.A.54 TOP: Graphing Logarithmic Functions

236 ANS: 3

$$\frac{-b}{a} = \frac{-6}{2} = -3. \quad \frac{c}{a} = \frac{4}{2} = 2$$

PTS: 2 REF: 011121a2 STA: A2.A.21 TOP: Roots of Quadratics
KEY: basic

237 ANS: 3

PTS: 2 REF: 081007a2 STA: A2.A.64
TOP: Using Inverse Trigonometric Functions KEY: basic

238 ANS: 3

$$S = \frac{-b}{a} = \frac{-(-3)}{4} = \frac{3}{4}. \quad P = \frac{c}{a} = \frac{-8}{4} = -2$$

PTS: 2 REF: fall0912a2 STA: A2.A.21 TOP: Roots of Quadratics
KEY: basic

239 ANS: 2

$$x^3 + x^2 - 2x = 0$$

$$x(x^2 + x - 2) = 0$$

$$x(x+2)(x-1) = 0$$

$$x = 0, -2, 1$$

PTS: 2 REF: 011103a2 STA: A2.A.26 TOP: Solving Polynomial Equations

240 ANS: 1

PTS: 2 REF: 061025a2 STA: A2.A.34
TOP: Sigma Notation

241 ANS: 4

PTS: 2 REF: 061120a2 STA: A2.A.19
TOP: Properties of Logarithms KEY: splitting logs

242 ANS: 2

$$\frac{x^{-1} - 1}{x - 1} = \frac{\frac{1}{x} - 1}{x - 1} = \frac{\frac{1-x}{x}}{x-1} = \frac{\frac{-(x-1)}{x}}{x-1} = -\frac{1}{x}$$

PTS: 2 REF: 081018a2 STA: A2.A.9 TOP: Negative Exponents

243 ANS: 4

$$2\log_4(5x) = 3$$

$$\log_4(5x) = \frac{3}{2}$$

$$5x = 4^{\frac{3}{2}}$$

$$5x = 8$$

$$x = \frac{8}{5}$$

PTS: 2

REF: fall0921a2

STA: A2.A.28

TOP: Logarithmic Equations

KEY: advanced

244 ANS: 4

$$b^2 - 4ac = 3^2 - 4(9)(-4) = 9 + 144 = 153$$

PTS: 2

REF: 081016a2

STA: A2.A.2

TOP: Using the Discriminant

KEY: determine nature of roots given equation

245 ANS: 1

$$13^2 = 15^2 + 14^2 - 2(15)(14)\cos C$$

$$169 = 421 - 420\cos C$$

$$-252 = -420\cos C$$

$$\frac{252}{420} = \cos C$$

$$53 \approx C$$

PTS: 2

REF: 061110a2

STA: A2.A.73

TOP: Law of Cosines

KEY: find angle

246 ANS: 2

$$\frac{10}{\sin 35} = \frac{13}{\sin B} \quad \cdot \quad 35 + 48 < 180$$

$$B \approx 48, 132 \quad 35 + 132 < 180$$

PTS: 2

REF: 011113a2

STA: A2.A.75

TOP: Law of Sines - The Ambiguous Case

247 ANS: 3

$$x = 5^4 = 625$$

PTS: 2

REF: 061106a2

STA: A2.A.28

TOP: Logarithmic Equations

KEY: basic

248 ANS: 3

PTS: 2

REF: 011104a2

STA: A2.A.64

TOP: Using Inverse Trigonometric Functions

KEY: unit circle

249 ANS: 3

PTS: 2

REF: 061127a2

STA: A2.S.6

TOP: Regression

250 ANS: 4

$$4ab\sqrt{2b} - 3a\sqrt{9b^2}\sqrt{2b} + 7ab\sqrt{6b} = 4ab\sqrt{2b} - 9ab\sqrt{2b} + 7ab\sqrt{6b} = -5ab\sqrt{2b} + 7ab\sqrt{6b}$$

PTS: 2 REF: fall0918a2 STA: A2.A.14 TOP: Operations with Radicals
KEY: with variables | index = 2

251 ANS: 3

$$68\% \times 50 = 34$$

PTS: 2 REF: 081013a2 STA: A2.S.5 TOP: Normal Distributions
KEY: predict

252 ANS: 1

$$\cos^2\theta - \cos 2\theta = \cos^2\theta - (\cos^2\theta - \sin^2\theta) = \sin^2\theta$$

PTS: 2 REF: 061024a2 STA: A2.A.77 TOP: Double Angle Identities
KEY: simplifying

253 ANS: 4

TOP: Analysis of Data

PTS: 2 REF: 061101a2 STA: A2.S.1

254 ANS: 3

TOP: Angle Sum and Difference Identities

PTS: 2 REF: fall0910a2 STA: A2.A.76
KEY: simplifying

255 ANS: 2

$$\frac{2\pi}{b} = \frac{2\pi}{3}$$

PTS: 2 REF: 061111a2 STA: A2.A.69
TOP: Properties of Graphs of Trigonometric Functions KEY: period

256 ANS: 1

TOP: Differentiating Permutations and Combinations

PTS: 2 REF: 011117a2 STA: A2.S.9

257 ANS: 4

$$7^2 = 3^2 + 5^2 - 2(3)(5)\cos A$$

$$49 = 34 - 30\cos A$$

$$15 = -30\cos A$$

$$-\frac{1}{2} = \cos A$$

$$120 = A$$

PTS: 2 REF: 081017a2 STA: A2.A.73 TOP: Law of Cosines
KEY: angle, without calculator

258 ANS: 4

TOP: Evaluating Logarithmic Expressions

PTS: 2 REF: 011124a2 STA: A2.A.18

259 ANS: 4

$$(3 + \sqrt{5})(3 - \sqrt{5}) = 9 - \sqrt{25} = 4$$

PTS: 2 REF: 081001a2 STA: A2.N.4 TOP: Operations with Irrational Expressions
KEY: without variables | index = 2

- 260 ANS: 4 PTS: 2 REF: 061124a2 STA: A2.S.3
TOP: Average Known with Missing Data
- 261 ANS: 2 PTS: 2 REF: 011126a2 STA: A2.A.49
TOP: Equations of Circles
- 262 ANS: 4 PTS: 2 REF: 061026a2 STA: A2.A.29
TOP: Sequences
- 263 ANS: 3 PTS: 2 REF: 081027a2 STA: A2.A.44
TOP: Inverse of Functions KEY: equations
- 264 ANS: 4

$$\frac{2x+4}{\sqrt{x+2}} \cdot \frac{\sqrt{x+2}}{\sqrt{x+2}} = \frac{2(x+2)\sqrt{x+2}}{x+2} = 2\sqrt{x+2}$$

- PTS: 2 REF: 011122a2 STA: A2.A.15 TOP: Rationalizing Denominators
KEY: index = 2

265 ANS: 2

$$\frac{11\pi}{12} \cdot \frac{180}{\pi} = 165$$

- PTS: 2 REF: 061002a2 STA: A2.M.2 TOP: Radian Measure
KEY: degrees

266 ANS: 3

$$f(4) = \frac{1}{2}(4) - 3 = -1. \quad g(-1) = 2(-1) + 5 = 3$$

- PTS: 2 REF: fall0902a2 STA: A2.A.42 TOP: Compositions of Functions
KEY: numbers

267 ANS: 3 PTS: 2 REF: fall0923a2 STA: A2.A.39
TOP: Domain and Range KEY: real domain

268 ANS: 1


$$4a + 6 = 4a - 10. \quad 4a + 6 = -4a + 10. \quad \left| 4\left(\frac{1}{2}\right) + 6 \right| - 4\left(\frac{1}{2}\right) = -10$$

$$6 \neq -10 \quad 8a = 4 \quad 8 - 2 \neq -10$$

$$a = \frac{4}{8} = \frac{1}{2}$$

- PTS: 2 REF: 011106a2 STA: A2.A.1 TOP: Absolute Value Equations

269 ANS: 2



$$\frac{\frac{\pi}{3} + \frac{\pi}{3}}{2\pi} = \frac{\frac{2\pi}{3}}{2\pi} = \frac{1}{3}$$

PTS: 2

REF: 011108a2

STA: A2.S.13

TOP: Geometric Probability

270 ANS: 3

$$27r^{4-1} = 64$$

$$r^3 = \frac{64}{27}$$

$$r = \frac{4}{3}$$

PTS: 2

REF: 081025a2

STA: A2.A.31

TOP: Sequences

271 ANS: 2

$$\cos(-305^\circ + 360^\circ) = \cos(55^\circ)$$

PTS: 2

REF: 061104a2

STA: A2.A.57

TOP: Reference Angles

272 ANS: 2

$$K = \frac{1}{2}(10)(18)\sin 120 = 45\sqrt{3} \approx 78$$

PTS: 2

REF: fall0907a2

STA: A2.A.74

TOP: Using Trigonometry to Find Area

KEY: basic

273 ANS: 3

```


Plot1 Plot2 Plot3
Y1=1/cos(X)
Y2=
Y3=
Y4=
Y5=
Y6=
Y7=

```

```

WINDOW
Xmin=0
Xmax=6.2831853...
Xscl=1.5707963...
Ymin=-2
Ymax=2
Yscl=1
Xres=1

```



PTS: 2

REF: 061020a2

STA: A2.A.71

TOP: Graphing Trigonometric Functions

274 ANS: 1

$${}_5C_3(3x)^2(-2)^3 = 10 \cdot 9x^2 \cdot -8 = -720x^2$$

PTS: 2

REF: fall0919a2

STA: A2.A.36

TOP: Binomial Expansions

275 ANS: 3

$$\frac{3^{-2}}{(-2)^{-3}} = \frac{\frac{1}{9}}{-\frac{1}{8}} = -\frac{8}{9}$$

PTS: 2 REF: 061003a2 STA: A2.N.1 TOP: Negative and Fractional Exponents

276 ANS: 1

$${}_{10}C_4 = 210$$

PTS: 2 REF: 061113a2 STA: A2.S.11 TOP: Combinations

277 ANS: 4

(4) fails the horizontal line test. Not every element of the range corresponds to only one element of the domain.

PTS: 2 REF: fall0906a2 STA: A2.A.43 TOP: Defining Functions


278 ANS: 1

$$-420 \left(\frac{\pi}{180} \right) = -\frac{7\pi}{3}$$

PTS: 2 REF: 081002a2 STA: A2.M.2 TOP: Radian Measure

KEY: radians

279 ANS: 2



A calculator screen showing the calculation of the tangent of an angle. The input is $\tan(126^\circ 43')$ and the output is -1.340788784 .

PTS: 2 REF: 061115a2 STA: A2.A.66 TOP: Determining Trigonometric Functions

280 ANS: 1

$$\frac{\sqrt{3} + 5}{\sqrt{3} - 5} \cdot \frac{\sqrt{3} + 5}{\sqrt{3} + 5} = \frac{3 + 5\sqrt{3} + 5\sqrt{3} + 25}{3 - 25} = \frac{28 + 10\sqrt{3}}{-22} = -\frac{14 + 5\sqrt{3}}{11}$$

PTS: 2 REF: 061012a2 STA: A2.N.5 TOP: Rationalizing Denominators

281 ANS: 1

$${}_9C_3 a^6 (-4b)^3 = -5376a^6 b^3$$

PTS: 2 REF: 061126a2 STA: A2.A.36 TOP: Binomial Expansions

282 ANS: 4

TOP: Permutations

PTS: 2 REF: fall0925a2 STA: A2.S.10

283 ANS: 1

$$\sqrt[4]{16x^2y^7} = 16^{\frac{1}{4}} x^{\frac{2}{4}} y^{\frac{7}{4}} = 2x^{\frac{1}{2}} y^{\frac{7}{4}}$$

PTS: 2 REF: 061107a2 STA: A2.A.11 TOP: Radicals as Fractional Exponents

284 ANS: 3

Cofunctions tangent and cotangent are complementary

PTS: 2 REF: 061014a2 STA: A2.A.58 TOP: Cofunction Trigonometric Relationships

285 ANS: 3

(1) and (4) fail the horizontal line test and are not one-to-one. Not every element of the range corresponds to only one element of the domain. (2) fails the vertical line test and is not a function. Not every element of the domain corresponds to only one element of the range.

PTS: 2 REF: 081020a2 STA: A2.A.43 TOP: Defining Functions

286 ANS: 3

n	0	1	2	Σ
$n^2 + 2^n$	$0^2 + 2^0 = 1$	$1^2 + 2^1 = 3$	$2^2 + 2^2 = 8$	12

$$2 \times 12 = 24$$

PTS: 2 REF: fall0911a2 STA: A2.N.10 TOP: Sigma Notation

KEY: basic

287 ANS: 2

$$x^2 - 2x + y^2 + 6y = -3$$

$$x^2 - 2x + 1 + y^2 + 6y + 9 = -3 + 1 + 9$$

$$(x - 1)^2 + (y + 3)^2 = 7$$

PTS: 2 REF: 061016a2 STA: A2.A.47 TOP: Equations of Circles

288 ANS: 4 PTS: 2 REF: 011101a2 STA: A2.A.38

TOP: Defining Functions KEY: graphs

289 ANS: 1

$$2 \log x - (3 \log y + \log z) = \log x^2 - \log y^3 - \log z = \log \frac{x^2}{y^3 z}$$

PTS: 2 REF: 061010a2 STA: A2.A.19 TOP: Properties of Logarithms

290 ANS: 2 PTS: 2 REF: 061122a2 STA: A2.A.24

TOP: Completing the Square

291 ANS: 2

$${}_{15}C_8 = 6,435$$

PTS: 2 REF: 081012a2 STA: A2.S.11 TOP: Combinations

292 ANS: 3 PTS: 2 REF: fall0913a2 STA: A2.A.65

TOP: Graphing Trigonometric Functions

293 ANS: 3 PTS: 2 REF: 061001a2 STA: A2.A.30

TOP: Sequences

294 ANS: 3 PTS: 2 REF: 061022a2 STA: A2.A.63

TOP: Domain and Range

295 ANS: 1 PTS: 2 REF: 011112a2 STA: A2.A.64

TOP: Using Inverse Trigonometric Functions KEY: advanced

296 ANS: 4

$$6x - x^3 - x^2 = -x(x^2 + x - 6) = -x(x + 3)(x - 2)$$

PTS: 2

REF: fall0917a2

STA: A2.A.7

TOP: Factoring Polynomials

KEY: single variable

297 ANS: 2

PTS: 2

REF: 061021a2

STA: A2.S.8

TOP: Correlation Coefficient

Algebra 2/Trigonometry 2 Point Regents Exam Questions Answer Section

298 ANS:

$$\sec \theta \sin \theta \cot \theta = \frac{1}{\cos \theta} \cdot \sin \theta \cdot \frac{\cos \theta}{\sin \theta} = 1$$

PTS: 2 REF: 011428a2 STA: A2.A.58 TOP: Reciprocal Trigonometric Relationships

299 ANS:

$$\cos \theta \cdot \frac{1}{\cos \theta} - \cos^2 \theta = 1 - \cos^2 \theta = \sin^2 \theta$$

PTS: 2 REF: 061230a2 STA: A2.A.58 TOP: Reciprocal Trigonometric Relationships

300 ANS:

$$3 - 2x \geq 7 \quad \text{or} \quad 3 - 2x \leq -7$$

$$-2x \geq 4 \quad -2x \leq -10$$

$$x \leq -2 \quad x \geq 5$$

PTS: 2 REF: 011334a2 STA: A2.A.1 TOP: Absolute Value Inequalities

KEY: graph

301 ANS:

$${}_7C_3 \left(\frac{1}{4} \right)^3 \left(\frac{3}{4} \right)^4 = 35 \left(\frac{1}{64} \right) \left(\frac{81}{256} \right) = \frac{2835}{16384} \approx 0.173$$

PTS: 2 REF: 061335a2 STA: A2.S.15 TOP: Binomial Probability

KEY: exactly

302 ANS:

$$5 \csc \theta = 8$$

$$\csc \theta = \frac{8}{5}$$

$$\sin \theta = \frac{5}{8}$$

$$\theta \approx 141$$

PTS: 2 REF: 061332a2 STA: A2.A.68 TOP: Trigonometric Equations

KEY: reciprocal functions

303 ANS:

$$x^2 - 6x - 27 = 0, \quad \frac{-b}{a} = 6, \quad \frac{c}{a} = -27. \quad \text{If } a = 1 \text{ then } b = -6 \text{ and } c = -27$$

PTS: 4 REF: 061130a2 STA: A2.A.21 TOP: Roots of Quadratics

KEY: basic

304 ANS:

$$\frac{\sin^2 A}{\cos^2 A} + \frac{\cos^2 A}{\cos^2 A} = \frac{1}{\cos^2 A}$$

$$\tan^2 A + 1 = \sec^2 A$$


PTS: 2 REF: 011135a2 STA: A2.A.67 TOP: Proving Trigonometric Identities

305 ANS:

$$y = 180.377(0.954)^x$$

PTS: 2 REF: 061231a2 STA: A2.S.7 TOP: Exponential Regression

306 ANS:



A calculator display showing the calculation of $3 \times \frac{180}{\pi}$ in degrees, minutes, and seconds. The display shows $(3 * \frac{180}{\pi}) \rightarrow \text{DMS}$ and the result $171^\circ 53' 14.419''$.

$$3 \times \frac{180}{\pi} \approx 171.89^\circ \approx 171^\circ 53'$$

PTS: 2 REF: 011135a2 STA: A2.M.2 TOP: Radian Measure
KEY: degrees

307 ANS:

$$2.5 \cdot \frac{180}{\pi} \approx 143^\circ 14'$$

PTS: 2 REF: 061431a2 STA: A2.M.2 TOP: Radian Measure
KEY: degrees

308 ANS:

$$D: -5 \leq x \leq 8. \quad R: -3 \leq y \leq 2$$

PTS: 2 REF: 011132a2 STA: A2.A.51 TOP: Domain and Range

309 ANS:

$$\frac{\sqrt{13}}{2} \cdot \sin \theta = \frac{y}{\sqrt{x^2 + y^2}} = \frac{2}{\sqrt{(-3)^2 + 2^2}} = \frac{2}{\sqrt{13}} \cdot \csc \theta = \frac{\sqrt{13}}{2}$$

PTS: 2 REF: fall0933a2 STA: A2.A.62 TOP: Determining Trigonometric Functions

310 ANS:

Controlled experiment because Howard is comparing the results obtained from an experimental sample against a control sample.

PTS: 2 REF: 081030a2 STA: A2.S.1 TOP: Analysis of Data

311 ANS:

$$y = 10.596(1.586)^x$$

PTS: 2 REF: 081031a2 STA: A2.S.7 TOP: Exponential Regression

312 ANS:

$$16^{2x+3} = 64^{x+2}$$

$$(4^2)^{2x+3} = (4^3)^{x+2}$$

$$4x + 6 = 3x + 6$$

$$x = 0$$

PTS: 2 REF: 011128a2 STA: A2.A.27 TOP: Exponential Equations
KEY: common base not shown

313 ANS:

$$7. f(-3) = (-3)^2 - 6 = 3. g(x) = 2^3 - 1 = 7.$$

PTS: 2 REF: 061135a2 STA: A2.A.42 TOP: Compositions of Functions
KEY: numbers

314 ANS:

$$7. 4 - \sqrt{2x-5} = 1$$

$$-\sqrt{2x-5} = -3$$

$$2x - 5 = 9$$

$$2x = 14$$

$$x = 7$$

PTS: 2 REF: 011229a2 STA: A2.A.22 TOP: Solving Radicals
KEY: basic

315 ANS:

$$12t^8 - 75t^4 = 3t^4(4t^4 - 25) = 3t^4(2t^2 + 5)(2t^2 - 5)$$

PTS: 2 REF: 061133a2 STA: A2.A.7
TOP: Factoring the Difference of Perfect Squares KEY: binomial

316 ANS:

$$a_1 = 3. a_2 = 2(3) - 1 = 5. a_3 = 2(5) - 1 = 9.$$

PTS: 2 REF: 061233a2 STA: A2.A.33 TOP: Recursive Sequences

317 ANS:

$$2x - 1 = 27^{\frac{4}{3}}$$

$$2x - 1 = 81$$

$$2x = 82$$

$$x = 41$$

PTS: 2 REF: 061329a2 STA: A2.A.28 TOP: Logarithmic Equations

KEY: advanced

318 ANS:

$$2.5 \cdot \frac{180}{\pi} \approx 143.2^\circ$$

PTS: 2 REF: 011129a2 STA: A2.M.2 TOP: Radian Measure

KEY: degrees

319 ANS:

$$3x^2 - 11x + 6 = 0. \text{ Sum } \frac{-b}{a} = \frac{11}{3}. \text{ Product } \frac{c}{a} = \frac{6}{3} = 2$$

PTS: 2 REF: 011329a2 STA: A2.A.20 TOP: Roots of Quadratics

320 ANS:

$$y = 0.488(1.116)^x$$

PTS: 2 REF: 061429a2 STA: A2.S.7 TOP: Exponential Regression

321 ANS:

$$\text{no solution. } \frac{4x}{x-3} = 2 + \frac{12}{x-3}$$

$$\frac{4x-12}{x-3} = 2$$

$$\frac{4(x-3)}{x-3} = 2$$

$$4 \neq 2$$

PTS: 2 REF: fall0930a2 STA: A2.A.23 TOP: Solving Rationals

KEY: rational solutions

322 ANS:

$$x < -1 \text{ or } x > 5. \quad x^2 - 4x - 5 > 0. \quad x - 5 > 0 \text{ and } x + 1 > 0 \text{ or } x - 5 < 0 \text{ and } x + 1 < 0$$

$$(x-5)(x+1) > 0 \quad x > 5 \text{ and } x > -1 \quad x < 5 \text{ and } x < -1$$

$$x > 5$$

$$x < -1$$

PTS: 2 REF: 011228a2 STA: A2.A.4 TOP: Quadratic Inequalities

KEY: one variable

323 ANS:

$$A = 750e^{(0.03)(8)} \approx 953$$

PTS: 2 REF: 061229a2 STA: A2.A.12 TOP: Evaluating Exponential Expressions

324 ANS:

$$\frac{31-19}{7-4} = \frac{12}{3} = 4 \quad x + (4-1)4 = 19 \quad a_n = 7 + (n-1)4$$

$$x + 12 = 19$$

$$x = 7$$

PTS: 2 REF: 011434a2 STA: A2.A.29 TOP: Sequences

325 ANS:

$$r = \sqrt{2^2 + 3^2} = \sqrt{13}. \quad (x+5)^2 + (y-2)^2 = 13$$

PTS: 2 REF: 011234a2 STA: A2.A.49 TOP: Writing Equations of Circles

326 ANS:

$$a + 15 + 2a = 90$$

$$3a + 15 = 90$$

$$3a = 75$$

$$a = 25$$

PTS: 2 REF: 011330a2 STA: A2.A.58 TOP: Cofunction Trigonometric Relationships

327 ANS:

$$g(10) = \left(a(10)\sqrt{1-x} \right)^2 = 100a^2(-9) = -900a^2$$

PTS: 2 REF: 061333a2 STA: A2.A.41 TOP: Functional Notation

328 ANS:

$$i^{13} + i^{18} + i^{31} + n = 0$$


$$i + (-1) - i + n = 0$$

$$-1 + n = 0$$

$$n = 1$$

PTS: 2 REF: 061228a2 STA: A2.N.7 TOP: Imaginary Numbers

329 ANS:

 $y = 0$

PTS: 2 REF: 061031a2 STA: A2.A.53 TOP: Graphing Exponential Functions

330 ANS:

$${}_{25}C_{20} = 53,130$$

PTS: 2 REF: 011232a2 STA: A2.S.11 TOP: Combinations

331 ANS:

$$\sec x = \sqrt{2}$$

$$\cos x = \frac{1}{\sqrt{2}}$$

$$\cos x = \frac{\sqrt{2}}{2}$$

$$x = 45^\circ, 315^\circ$$

PTS: 2 REF: 061434a2 STA: A2.A.68 TOP: Trigonometric Equations
KEY: reciprocal functions

332 ANS:

$$\frac{2\sqrt{3}}{3}. \text{ If } \sin 60 = \frac{\sqrt{3}}{2}, \text{ then } \csc 60 = \frac{2}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$$

PTS: 2 REF: 011235a2 STA: A2.A.59 TOP: Reciprocal Trigonometric Relationships

333 ANS:

$$\frac{\frac{1}{2} - \frac{4}{d}}{\frac{1}{d} + \frac{3}{2d}} = \frac{\frac{d-8}{2d}}{\frac{2d+3d}{2d^2}} = \frac{d-8}{2d} \times \frac{2d^2}{5d} = \frac{d-8}{5}$$

PTS: 2 REF: 061035a2 STA: A2.A.17 TOP: Complex Fractions

334 ANS:

$$\text{Sum } \frac{-b}{a} = -\frac{11}{5}, \text{ Product } \frac{c}{a} = -\frac{3}{5}$$

PTS: 2

REF: 061030a2

STA: A2.A.20

TOP: Roots of Quadratics

335 ANS:

$$4xi + 5yi^8 + 6xi^3 + 2yi^4 = 4xi + 5y - 6xi + 2y = 7y - 2xi$$


PTS: 2

REF: 011433a2

STA: A2.N.7

TOP: Imaginary Numbers

336 ANS:



41,040.

PTS: 2

REF: fall0935a2

STA: A2.S.12

TOP: Sample Space

337 ANS:

$$x(x + 3) = 10$$

$$x^2 + 3x - 10 = 0$$

$$(x + 5)(x - 2) = 0$$

$$x = -5, 2$$

PTS: 2

REF: 011431a2

STA: A2.A.3

TOP: Quadratic-Linear Systems

KEY: equations

338 ANS:

$y = -3 \sin 2x$. The period of the function is π , the amplitude is 3 and it is reflected over the x -axis.

PTS: 2

REF: 061235a2

STA: A2.A.72

TOP: Identifying the Equation of a Trigonometric Graph

339 ANS:

$$-4x + 5 < 13 \quad -4x + 5 > -13 \quad -2 < x < 4.5$$

$$-4x < 8 \quad -4x > -18$$

$$x > -2 \quad x < 4.5$$

PTS: 2

REF: 011432a2

STA: A2.A.1

TOP: Absolute Value Inequalities

340 ANS:

$$e^{3 \ln 2} = e^{\ln 2^3} = e^{\ln 8} = 8$$

PTS: 2

REF: 061131a2

STA: A2.A.12

TOP: Evaluating Exponential Expressions

341 ANS:

$$5\sqrt{3x^3} - 2\sqrt{27x^3} = 5\sqrt{x^2} \sqrt{3x} - 2\sqrt{9x^2} \sqrt{3x} = 5x\sqrt{3x} - 6x\sqrt{3x} = -x\sqrt{3x}$$

PTS: 2 REF: 061032a2 STA: A2.N.2 TOP: Operations with Radicals

342 ANS:

$$-\frac{a^2b^3}{4}$$

PTS: 2 REF: 011231a2 STA: A2.A.13 TOP: Simplifying Radicals
KEY: index > 2

343 ANS:

$$\frac{5(3+\sqrt{2})}{7} \cdot \frac{5}{3-\sqrt{2}} \times \frac{3+\sqrt{2}}{3+\sqrt{2}} = \frac{5(3+\sqrt{2})}{9-2} = \frac{5(3+\sqrt{2})}{7}$$

PTS: 2 REF: fall0928a2 STA: A2.N.5 TOP: Rationalizing Denominators

344 ANS:

$$b^2 - 4ac = 0$$

$$k^2 - 4(1)(4) = 0$$


$$k^2 - 16 = 0$$

$$(k+4)(k-4) = 0$$

$$k = \pm 4$$

PTS: 2 REF: 061028a2 STA: A2.A.2 TOP: Using the Discriminant
KEY: determine equation given nature of roots

345 ANS:



PTS: 2 REF: 011234a2 STA: A2.A.53 TOP: Graphing Exponential Functions

346 ANS:

$$39,916,800 \cdot \frac{{}^{12}P_{12}}{3! \cdot 2!} = \frac{479,001,600}{12} = 39,916,800$$

PTS: 2 REF: 081035a2 STA: A2.S.10 TOP: Permutations

347 ANS:

$$K = ab \sin C = 6 \cdot 6 \sin 50 \approx 27.6$$

PTS: 2

REF: 011429a2

STA: A2.A.74

TOP: Using Trigonometry to Find Area

KEY: Parallelograms

348 ANS:

$$10ax^2 - 23ax - 5a = a(10x^2 - 23x - 5) = a(5x + 1)(2x - 5)$$

PTS: 2

REF: 081028a2

STA: A2.A.7

TOP: Factoring Polynomials

KEY: multiple variables

349 ANS:

$$\text{Sum } \frac{-b}{a} = -\frac{1}{12}, \text{ Product } \frac{c}{a} = -\frac{1}{2}$$


PTS: 2

REF: 061328a2

STA: A2.A.20

TOP: Roots of Quadratics

350 ANS:



PTS: 2


REF: 061033a2

STA: A2.A.60

TOP: Unit Circle

351 ANS:

$$197^\circ 40'. \quad 3.45 \times \frac{180}{\pi} \approx 197^\circ 40'.$$



PTS: 2

REF: fall0931a2

STA: A2.M.2

TOP: Radian Measure

KEY: degrees

352 ANS:

$$\frac{\sqrt{3}}{2} \times \frac{\sqrt{2}}{2} = \frac{\sqrt{6}}{4}$$

PTS: 2 REF: 061331a2 STA: A2.A.56 TOP: Determining Trigonometric Functions
KEY: degrees, common angles

353 ANS:

-3, -5, -8, -12

PTS: 2 REF: fall0934a2 STA: A2.A.33 TOP: Recursive Sequences

354 ANS:

$$45, 225 \quad 2 \tan C - 3 = 3 \tan C - 4$$

$$1 = \tan C$$

$$\tan^{-1} 1 = C$$

$$C = 45, 225$$

PTS: 2 REF: 081032a2 STA: A2.A.68 TOP: Trigonometric Equations
KEY: basic

355 ANS:

$$(x + 5)^2 + (y - 3)^2 = 32$$

PTS: 2 REF: 081033a2 STA: A2.A.49 TOP: Writing Equations of Circles

356 ANS:

Less than 60 inches is below 1.5 standard deviations from the mean. $0.067 \cdot 450 \approx 30$

PTS: 2 REF: 061428a2 STA: A2.S.5 TOP: Normal Distributions
KEY: predict

357 ANS:

$$(5x - 1)^{\frac{1}{3}} = 4$$

$$5x - 1 = 64$$

$$5x = 65$$

$$x = 13$$

PTS: 2 REF: 061433a2 STA: A2.A.28 TOP: Logarithmic Equations
KEY: advanced

358 ANS:

$$(x + 3)^2 + (y - 4)^2 = 25$$

PTS: 2 REF: fall0929a2 STA: A2.A.49 TOP: Writing Equations of Circles

359 ANS:

$$83^\circ 50' \cdot \frac{\pi}{180} \approx 1.463 \text{ radians } s = \theta r = 1.463 \cdot 12 \approx 17.6$$

PTS: 2 REF: 011435a2 STA: A2.A.61 TOP: Arc Length

KEY: arc length

360 ANS:

$$\frac{2 \pm \sqrt{(-2)^2 - 4(6)(-3)}}{2(6)} = \frac{2 \pm \sqrt{76}}{12} = \frac{2 \pm \sqrt{4} \sqrt{19}}{12} = \frac{2 \pm 2\sqrt{19}}{12} = \frac{1 \pm \sqrt{19}}{6}$$

PTS: 2 REF: 011332a2 STA: A2.A.25 TOP: Quadratics with Irrational Solutions

361 ANS:

Ordered, the heights are 71, 71, 72, 74, 74, 75, 78, 79, 79, 83. $Q_1 = 72$ and $Q_3 = 79$. $79 - 72 = 7$.

PTS: 2 REF: 011331a2 STA: A2.S.4 TOP: Dispersion

KEY: range, quartiles, interquartile range, variance

362 ANS:

$$\sum_{n=1}^{15} 7n$$

PTS: 2 REF: 081029a2 STA: A2.A.34 TOP: Sigma Notation

363 ANS:

$$12 \cdot 6 = 9w$$

$$8 = w$$

PTS: 2 REF: 011130a2 STA: A2.A.5 TOP: Inverse Variation

364 ANS:

$$a_n = 9n - 4 \quad \cdot \quad S_n = \frac{20(5 + 176)}{2} = 1810$$

$$a_1 = 9(1) - 4 = 5$$

$$a_{20} = 9(20) - 4 = 176$$

PTS: 2 REF: 011328a2 STA: A2.A.35 TOP: Summations

KEY: arithmetic

365 ANS:

$$\frac{\sqrt{108x^5y^8}}{\sqrt{6xy^5}} = \sqrt{18x^4y^3} = 3x^2y\sqrt{2y}$$

PTS: 2 REF: 011133a2 STA: A2.A.14 TOP: Operations with Radicals

KEY: with variables | index = 2

366 ANS:

$$230. 10 + (1^3 - 1) + (2^3 - 1) + (3^3 - 1) + (4^3 - 1) + (5^3 - 1) = 10 + 0 + 7 + 26 + 63 + 124 = 230$$

PTS: 2 REF: 011131a2 STA: A2.N.10 TOP: Sigma Notation
KEY: basic

367 ANS:

no. over 20 is more than 1 standard deviation above the mean. $0.159 \cdot 82 \approx 13.038$

PTS: 2 REF: 061129a2 STA: A2.S.5 TOP: Normal Distributions
KEY: predict

368 ANS:

$$(x + yi)(x - yi) = x^2 - y^2i^2 = x^2 + y^2$$

PTS: 2 REF: 061432a2 STA: A2.N.8 TOP: Conjugates of Complex Numbers

369 ANS:

7.4


PTS: 2 REF: 061029a2 STA: A2.S.4 TOP: Dispersion
KEY: basic, group frequency distributions

370 ANS:

$$\frac{\cot x \sin x}{\sec x} = \frac{\frac{\cos x}{\sin x} \sin x}{\frac{1}{\cos x}} = \cos^2 x$$

PTS: 2 REF: 061334a2 STA: A2.A.58 TOP: Reciprocal Trigonometric Relationships

371 ANS:



PTS: 2 REF: 061435a2 STA: A2.A.46
TOP: Transformations with Functions and Relations

372 ANS:

68% of the students are within one standard deviation of the mean. 16% of the students are more than one standard deviation above the mean.

PTS: 2 REF: 011134a2 STA: A2.S.5 TOP: Normal Distributions
KEY: percent

373 ANS:

$$2x - 3 > 5 \text{ or } 2x - 3 < -5$$

$$2x > 8 \quad 2x < -2$$

$$x > 4 \quad x < -1$$

PTS: 2

REF: 061430a2

STA: A2.A.1

TOP: Absolute Value Inequalities

374 ANS:

$$\frac{12x^2}{y^9} \cdot \frac{3x^{-4}y^5}{(2x^3y^{-7})^{-2}} = \frac{3y^5(2x^3y^{-7})^2}{x^4} = \frac{3y^5(4x^6y^{-14})}{x^4} = \frac{12x^6y^{-9}}{x^4} = \frac{12x^2}{y^9}$$

PTS: 2

REF: 061134a2

STA: A2.A.9

TOP: Negative Exponents

375 ANS:

$$y = x^2 - 6. \quad f^{-1}(x) \text{ is not a function.}$$

$$x = y^2 - 6$$

$$x + 6 = y^2$$

$$\pm\sqrt{x+6} = y$$

PTS: 2

REF: 061132a2

STA: A2.A.44

TOP: Inverse of Functions

KEY: equations

376 ANS:

$$216\left(\frac{\pi}{180}\right) \approx 3.8$$

PTS: 2

REF: 061232a2

STA: A2.M.2

TOP: Radian Measure

KEY: radians

377 ANS:

$$30700 = 50e^{3t}$$

$$614 = e^{3t}$$

$$\ln 614 = \ln e^{3t}$$

$$\ln 614 = 3t \ln e$$

$$\ln 614 = 3t$$

$$2.14 \approx t$$

PTS: 2

REF: 011333a2

STA: A2.A.6

TOP: Exponential Growth

378 ANS:

$$\frac{4}{9}x^2 - \frac{4}{3}x + 1. \quad \left(\frac{2}{3}x - 1\right)^2 = \left(\frac{2}{3}x - 1\right)\left(\frac{2}{3}x - 1\right) = \frac{4}{9}x^2 - \frac{2}{3}x - \frac{2}{3}x + 1 = \frac{4}{9}x^2 - \frac{4}{3}x + 1$$

PTS: 2

REF: 081034a2

STA: A2.N.3

TOP: Operations with Polynomials

379 ANS:

$$Q_1 = 3.5 \text{ and } Q_3 = 10.5. \quad 10.5 - 3.5 = 7.$$

PTS: 2 REF: 011430a2 STA: A2.S.4 TOP: Dispersion


KEY: range, quartiles, interquartile range, variance

380 ANS:

$$6y^3 - \frac{37}{10}y^2 - \frac{1}{5}y \cdot \left(\frac{1}{2}y^2 - \frac{1}{3}y \right) \left(12y + \frac{3}{5} \right) = 6y^3 + \frac{3}{10}y^2 - 4y^2 - \frac{1}{5}y = 6y^3 - \frac{37}{10}y^2 - \frac{1}{5}y$$

PTS: 2 REF: 061128a2 STA: A2.N.3 TOP: Operations with Polynomials

381 ANS:



-104.

PTS: 2 REF: 011230a2 STA: A2.N.10 TOP: Sigma Notation

KEY: basic


382 ANS:

$$K = ab \sin C = 18 \cdot 22 \sin 60 = 396 \frac{\sqrt{3}}{2} = 198\sqrt{3}$$

PTS: 2 REF: 061234a2 STA: A2.A.74 TOP: Using Trigonometry to Find Area

KEY: Parallelograms

383 ANS:



2,298.65.

PTS: 2 REF: fall0932a2 STA: A2.A.12 TOP: Evaluating Exponential Expressions

384 ANS:

$$K = ab \sin C = 24 \cdot 30 \sin 57 \approx 604$$

PTS: 2 REF: 061034a2 STA: A2.A.74 TOP: Using Trigonometry to Find Area

KEY: parallelograms

385 ANS:

$$\frac{{}_{10}P_{10}}{3! \cdot 3! \cdot 2!} = \frac{3,628,800}{72} = 50,400$$

PTS: 2 REF: 061330a2 STA: A2.S.10 TOP: Permutations

Algebra 2/Trigonometry 4 Point Regents Exam Questions Answer Section

386 ANS:

$$(x + 4)^2 = 17x - 4$$

$$x^2 + 8x + 16 = 17x - 4$$

$$x^2 - 9x + 20 = 0$$

$$(x - 4)(x - 5) = 0$$

$$x = 4, 5$$

PTS: 4

REF: 011336a2

STA: A2.A.28

TOP: Logarithmic Equations

KEY: basic

387 ANS:

$$\frac{51}{243} \cdot {}_5C_3 \left(\frac{1}{3}\right)^3 \left(\frac{2}{3}\right)^2 = \frac{40}{243}$$

$${}_5C_4 \left(\frac{1}{3}\right)^4 \left(\frac{2}{3}\right)^1 = \frac{10}{243}$$

$${}_5C_3 \left(\frac{1}{3}\right)^5 \left(\frac{2}{3}\right)^0 = \frac{1}{243}$$

PTS: 4

REF: 061138a2

STA: A2.S.15

TOP: Binomial Probability

KEY: at least or at most


388 ANS:

$$-3|6 - x| < -15$$

$$|6 - x| > 5$$

$$6 - x > 5 \text{ or } 6 - x < -5$$

$$1 > x \text{ or } 11 < x$$



PTS: 2

REF: 061137a2

STA: A2.A.1

TOP: Absolute Value Inequalities

KEY: graph

389 ANS:

$$y = 215.983(1.652)^x. \quad 215.983(1.652)^7 \approx 7250$$

PTS: 4

REF: 011337a2

STA: A2.S.7

TOP: Exponential Regression

390 ANS:

$$y = 27.2025(1.1509)^x. \quad y = 27.2025(1.1509)^{18} \approx 341$$

PTS: 4

REF: 011238a2

STA: A2.S.7

TOP: Exponential Regression

391 ANS:

$$0.468. {}_8C_6 \left(\frac{2}{3}\right)^6 \left(\frac{1}{3}\right)^2 \approx 0.27313. {}_8C_7 \left(\frac{2}{3}\right)^7 \left(\frac{1}{3}\right)^1 \approx 0.15607. {}_8C_8 \left(\frac{2}{3}\right)^8 \left(\frac{1}{3}\right)^0 \approx 0.03902.$$

PTS: 4 REF: 011138a2 STA: A2.S.15 TOP: Binomial Probability


KEY: at least or at most

392 ANS:

$$\begin{aligned} \pm \frac{3}{2}, -\frac{1}{2}. \quad 8x^3 + 4x^2 - 18x - 9 &= 0 \\ 4x^2(2x+1) - 9(2x+1) &= 0 \\ (4x^2 - 9)(2x+1) &= 0 \\ 4x^2 - 9 = 0 \text{ or } 2x+1 = 0 & \\ (2x+3)(2x-3) = 0 \quad x = -\frac{1}{2} & \\ x = \pm \frac{3}{2} & \end{aligned}$$

PTS: 4 REF: fall0937a2 STA: A2.A.26 TOP: Solving Polynomial Equations

393 ANS:



$$\begin{aligned} \frac{27}{\sin 75} = \frac{F_1}{\sin 60} \cdot \frac{27}{\sin 75} = \frac{F_2}{\sin 45} \cdot \\ F_1 \approx 24 \quad F_2 \approx 20 \end{aligned}$$

PTS: 4 REF: 061238a2 STA: A2.A.73 TOP: Vectors

394 ANS:

$$\begin{aligned} \frac{100}{\sin 32} = \frac{b}{\sin 105} \cdot \frac{100}{\sin 32} = \frac{a}{\sin 43} \cdot \\ b \approx 182.3 \quad a \approx 128.7 \end{aligned}$$

PTS: 4 REF: 011338a2 STA: A2.A.73 TOP: Law of Sines

KEY: basic

395 ANS:

$$\begin{aligned} \sin(45 + 30) &= \sin 45 \cos 30 + \cos 45 \sin 30 \\ &= \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2} + \frac{\sqrt{2}}{2} \cdot \frac{1}{2} = \frac{\sqrt{6}}{4} + \frac{\sqrt{2}}{4} = \frac{\sqrt{6} + \sqrt{2}}{4} \end{aligned}$$

PTS: 4 REF: 061136a2 STA: A2.A.76 TOP: Angle Sum and Difference Identities

KEY: evaluating

396 ANS:

$$y = 2.001x^{2.298}, 1,009. \quad y = 2.001(15)^{2.298} \approx 1009$$

PTS: 4 REF: fall0938a2 STA: A2.S.7 TOP: Power Regression

397 ANS:

$$\text{No. TENNESSEE: } \frac{{}_9P_9}{4! \cdot 2! \cdot 2!} = \frac{362,880}{96} = 3,780. \quad \text{VERMONT: } {}_7P_7 = 5,040$$

PTS: 4 REF: 061038a2 STA: A2.S.10 TOP: Permutations

398 ANS:

0, 60, 180, 300.

$$\sin 2\theta = \sin \theta$$

$$\sin 2\theta - \sin \theta = 0$$

$$2\sin \theta \cos \theta - \sin \theta = 0$$

$$\sin \theta (2\cos \theta - 1) = 0$$

$$\sin \theta = 0 \quad 2\cos \theta - 1 = 0$$

$$\theta = 0, 180 \quad \cos \theta = \frac{1}{2}$$

$$\theta = 60, 300$$

PTS: 4 REF: 061037a2 STA: A2.A.68 TOP: Trigonometric Equations
KEY: double angle identities

399 ANS:

$$26.2\%. \quad {}_{10}C_8 \cdot 0.65^8 \cdot 0.35^2 + {}_{10}C_9 \cdot 0.65^9 \cdot 0.35^1 + {}_{10}C_{10} \cdot 0.65^{10} \cdot 0.35^0 \approx 0.262$$

PTS: 4 REF: 081038a2 STA: A2.S.15 TOP: Binomial Probability
KEY: at least or at most

400 ANS:

$$\frac{15}{\sin 103} = \frac{a}{\sin 42} \cdot \frac{1}{2} (15)(10.3)\sin 35 \approx 44$$

$$a \approx 10.3$$

PTS: 4 REF: 061337a2 STA: A2.A.74 TOP: Using Trigonometry to Find Area
KEY: advanced

401 ANS:

$$0.167. \quad {}_{10}C_8 \cdot 0.6^8 \cdot 0.4^2 + {}_{10}C_9 \cdot 0.6^9 \cdot 0.4^1 + {}_{10}C_{10} \cdot 0.6^{10} \cdot 0.4^0 \approx 0.167$$

PTS: 4 REF: 061036a2 STA: A2.S.15 TOP: Binomial Probability
KEY: at least or at most

402 ANS:

$$\sigma_x = 14.9. \quad \bar{x} = 40. \quad \text{There are 8 scores between 25.1 and 54.9.}$$
PTS: 4 REF: 061237a2 STA: A2.S.4 TOP: Dispersion
KEY: advanced

403 ANS:

$$88. \frac{100}{\sin 33} = \frac{x}{\sin 32} \cdot \sin 66 \approx \frac{T}{97.3}$$

$$x \approx 97.3 \quad t \approx 88$$

PTS: 4 REF: 011236a2 STA: A2.A.73 TOP: Law of Sines
KEY: advanced

404 ANS:

$$\sqrt{27^2 + 32^2 - 2(27)(32)\cos 132} \approx 54$$

PTS: 4 REF: 011438a2 STA: A2.A.73 TOP: Law of Cosines
KEY: applied

405 ANS:

$$2\sin^2 x + 5\sin x - 3 = 0$$

$$(2\sin x - 1)(\sin x + 3) = 0$$

$$\sin x = \frac{1}{2}$$

$$x = \frac{\pi}{6}, \frac{5\pi}{6}$$

PTS: 4 REF: 011436a2 STA: A2.A.68 TOP: Trigonometric Equations
KEY: quadratics

406 ANS:

$$\frac{13}{x} = 10 - x \quad \cdot \quad x = \frac{10 \pm \sqrt{100 - 4(1)(13)}}{2(1)} = \frac{10 \pm \sqrt{48}}{2} = \frac{10 \pm 4\sqrt{3}}{2} = 5 \pm 2\sqrt{3}$$

$$13 = 10x - x^2$$

$$x^2 - 10x + 13 = 0$$

PTS: 4 REF: 061336a2 STA: A2.A.23 TOP: Solving Rationals
KEY: irrational and complex solutions

407 ANS:

$$32x^5 - 80x^4 + 80x^3 - 40x^2 + 10x - 1 \cdot {}_5C_0(2x)^5(-1)^0 = 32x^5 \cdot {}_5C_1(2x)^4(-1)^1 = -80x^4 \cdot {}_5C_2(2x)^3(-1)^2 = 80x^3 \cdot {}_5C_3(2x)^2(-1)^3 = -40x^2 \cdot {}_5C_4(2x)^1(-1)^4 = 10x \cdot {}_5C_5(2x)^0(-1)^5 = -1$$

PTS: 4 REF: 011136a2 STA: A2.A.36 TOP: Binomial Expansions

408 ANS:

$$\frac{1 + \frac{3}{x}}{1 - \frac{5}{x} - \frac{24}{x^2}} \cdot \frac{x^2}{x^2} = \frac{x^2 + 3x}{x^2 - 5x - 24} = \frac{x(x+3)}{(x-8)(x+3)} = \frac{x}{x-8}$$

PTS: 4 REF: 061436a2 STA: A2.A.17 TOP: Complex Fractions

409 ANS:

$${}_5C_4 \cdot 0.28^4 \cdot 0.72^1 + {}_5C_5 \cdot 0.28^5 \cdot 0.72^0 \approx 0.024$$

PTS: 4 REF: 011437a2 STA: A2.S.15 TOP: Binomial Probability

KEY: at least or at most

410 ANS:

$$\frac{23}{2} \quad \cos^2 B + \sin^2 B = 1 \quad \tan B = \frac{\sin B}{\cos B} = \frac{\frac{5}{\sqrt{41}}}{\frac{4}{\sqrt{41}}} = \frac{5}{4}$$

$$\cos^2 B + \left(\frac{5}{\sqrt{41}} \right)^2 = 1$$

$$\cos^2 B + \frac{25}{41} = \frac{41}{41}$$

$$\cos^2 B = \frac{16}{41}$$

$$\cos B = \frac{4}{\sqrt{41}}$$

$$\tan(A + B) = \frac{\frac{2}{3} + \frac{5}{4}}{1 - \left(\frac{2}{3} \right) \left(\frac{5}{4} \right)} = \frac{\frac{8+15}{12}}{\frac{12}{12} - \frac{10}{12}} = \frac{\frac{23}{12}}{\frac{2}{12}} = \frac{23}{2}$$

PTS: 4 REF: 081037a2 STA: A2.A.76 TOP: Angle Sum and Difference Identities

KEY: evaluating

411 ANS:

$$800. \quad x = 4^{2.5} = 32. \quad y^{-\frac{3}{2}} = 125 \quad \cdot \frac{x}{y} = \frac{32}{\frac{1}{25}} = 800$$

$$y = 125^{-\frac{2}{3}} = \frac{1}{25}$$

PTS: 4 REF: 011237a2 STA: A2.A.28 TOP: Logarithmic Equations

KEY: advanced

412 ANS:

$$x^3 + 5x^2 - 4x - 20 = 0$$

$$x^2(x+5) - 4(x+5) = 0$$

$$(x^2 - 4)(x+5) = 0$$

$$(x+2)(x-2)(x+5) = 0$$

$$x = \pm 2, -5$$

PTS: 4 REF: 061437a2 STA: A2.A.26 TOP: Solving Polynomial Equations

413 ANS:

$${}_5C_0 \cdot 0.57^0 \cdot 0.43^5 + {}_5C_1 \cdot 0.57^1 \cdot 0.43^4 + {}_5C_2 \cdot 0.57^2 \cdot 0.43^3 \approx 0.37$$

PTS: 4 REF: 061438a2 STA: A2.S.15 TOP: Binomial Probability

KEY: at least or at most

414 ANS:

$$\frac{-(x^2-4)}{(x+4)(x+3)} \times \frac{x+3}{2(x-2)} = \frac{-(x+2)(x-2)}{x+4} \times \frac{1}{2(x-2)} = \frac{-(x+2)}{2(x+4)}$$

PTS: 4 REF: 061236a2 STA: A2.A.16 TOP: Multiplication and Division of Rationals

KEY: division

415 ANS:

$$3 \pm \sqrt{7}. \quad 2x^2 - 12x + 4 = 0$$

$$x^2 - 6x + 2 = 0$$

$$x^2 - 6x = -2$$

$$x^2 - 6x + 9 = -2 + 9$$

$$(x-3)^2 = 7$$

$$x-3 = \pm\sqrt{7}$$

$$x = 3 \pm \sqrt{7}$$

PTS: 4 REF: fall0936a2 STA: A2.A.24 TOP: Completing the Square

416 ANS:

$\sigma_x \approx 6.2$. 6 scores are within a population standard deviation of the mean. $Q_3 - Q_1 = 41 - 37 = 4$

$\bar{x} \approx 38.2$

PTS: 4 REF: 061338a2 STA: A2.S.4 TOP: Dispersion

KEY: advanced

417 ANS:

$$\frac{1}{3} - \frac{1}{x+3} - \frac{2}{3-x} = \frac{4}{x^2-9}$$

$$\frac{1}{x+3} + \frac{2}{x-3} = \frac{4}{x^2-9}$$

$$\frac{x-3+2(x+3)}{(x+3)(x-3)} = \frac{4}{(x+3)(x-3)}$$

$$x-3+2x+6=4$$

$$3x=1$$

$$x = \frac{1}{3}$$

PTS: 4 REF: 081036a2 STA: A2.A.23 TOP: Solving Rationals

KEY: rational solutions

418 ANS:

$$\frac{12}{\sin 32} = \frac{10}{\sin B} \quad . \quad C \approx 180 - (32 + 26.2) \approx 121.8. \quad \frac{12}{\sin 32} = \frac{c}{\sin 121.8}$$
$$B = \sin^{-1} \frac{10 \sin 32}{12} \approx 26.2 \quad c = \frac{12 \sin 121.8}{\sin 32} \approx 19.2$$

PTS: 4

REF: 011137a2

STA: A2.A.73

TOP: Law of Sines

KEY: basic

Algebra 2/Trigonometry 6 Point Regents Exam Questions Answer Section

419 ANS:

$$81^{x^3+2x^2} = 27^{\frac{5x}{3}}$$

$$\left(3^4\right)^{x^3+2x^2} = \left(3^3\right)^{\frac{5x}{3}}$$

$$3^{4x^3+8x^2} = 3^{5x}$$

$$4x^3 + 8x^2 - 5x = 0$$

$$x(4x^2 + 8x - 5) = 0$$

$$x(2x - 1)(2x + 5) = 0$$

$$x = 0, \frac{1}{2}, -\frac{5}{2}$$

PTS: 6 REF: 061239a2 STA: A2.A.27 TOP: Exponential Equations
KEY: common base not shown

420 ANS:

$$\sqrt{x^2 + x - 1} = -4x + 3 \quad -4\left(\frac{2}{3}\right) + 3 \geq 0$$

$$x^2 + x - 1 = 16x^2 - 24x + 9 \quad \frac{1}{3} \geq 0$$

$$0 = 15x^2 - 25x + 10$$

$$0 = 3x^2 - 5x + 2 \quad -4(1) + 3 < 0$$

$$0 = (3x - 2)(x - 1) \quad 1 \text{ is extraneous}$$

$$x = \frac{2}{3}, x \neq 1$$

PTS: 6 REF: 011339a2 STA: A2.A.22 TOP: Solving Radicals
KEY: extraneous solutions

421 ANS:

$$\frac{-2(x^2 + 6)}{x^4} \cdot \frac{x^2(x - 3) + 6(x - 3)}{x^2 - 4x} \cdot \frac{2x - 4}{x^4 - 3x^3} \div \frac{x^2 + 2x - 8}{16 - x^2}$$

$$\frac{(x^2 + 6)(x - 3)}{x(x - 4)} \cdot \frac{2(x - 2)}{x^3(x - 3)} \cdot \frac{(4 + x)(4 - x)}{(x + 4)(x - 2)}$$

$$\frac{-2(x^2 + 6)}{x^4}$$

PTS: 6 REF: 011239a2 STA: A2.A.16 TOP: Multiplication and Division of Rationals
KEY: division

422 ANS:

$$x = -\frac{1}{3}, -1 \quad \log_{x+3} \frac{x^3+x-2}{x} = 2$$

$$\frac{x^3+x-2}{x} = (x+3)^2$$

$$\frac{x^3+x-2}{x} = x^2+6x+9$$

$$x^3+x-2 = x^3+6x^2+9x$$

$$0 = 6x^2+8x+2$$

$$0 = 3x^2+4x+1$$

$$0 = (3x+1)(x+1)$$

$$x = -\frac{1}{3}, -1$$

PTS: 6

REF: 081039a2

STA: A2.A.28

TOP: Logarithmic Equations

KEY: basic

423 ANS:

$$x^4+4x^3+4x^2+16x=0$$

$$x(x^3+4x^2+4x+16)=0$$

$$x(x^2(x+4)+4(x+4))=0$$

$$x(x^2+4)(x+4)=0$$

$$x = 0, \pm 2i, -4$$

PTS: 6

REF: 061339a2

STA: A2.A.26

TOP: Solving Polynomial Equations

424 ANS:

$$\left(-\frac{9}{2}, \frac{1}{2}\right) \text{ and } \left(\frac{1}{2}, \frac{11}{2}\right). \quad y = x + 5 \quad . \quad 4x^2 + 17x - 4 = x + 5$$

$$y = 4x^2 + 17x - 4 \quad 4x^2 + 16x - 9 = 0$$

$$(2x+9)(2x-1) = 0$$

$$x = -\frac{9}{2} \text{ and } x = \frac{1}{2}$$

$$y = -\frac{9}{2} + 5 = \frac{1}{2} \text{ and } y = \frac{1}{2} + 5 = \frac{11}{2}$$

PTS: 6

REF: 061139a2

STA: A2.A.3

TOP: Quadratic-Linear Systems

KEY: equations

425 ANS:

$$\log_{(x+3)}(2x+3)(x+5) = 2 \quad -6 \text{ is extraneous}$$

$$(x+3)^2 = (2x+3)(x+5)$$

$$x^2 + 6x + 9 = 2x^2 + 13x + 15$$

$$x^2 + 7x + 6 = 0$$

$$(x+6)(x+1) = 0$$

$$x = -1$$

PTS: 6 REF: 011439a2 STA: A2.A.28 TOP: Logarithmic Equations

KEY: applying properties of logarithms

426 ANS:

$$33. a = \sqrt{10^2 + 6^2 - 2(10)(6)\cos 80} \approx 10.7. \angle C \text{ is opposite the shortest side. } \frac{6}{\sin C} = \frac{10.7}{\sin 80}$$

$$C \approx 33$$

PTS: 6 REF: 061039a2 STA: A2.A.73 TOP: Law of Cosines

KEY: advanced

427 ANS:

$$\ln(T - T_0) = -kt + 4.718 \quad \ln(T - 68) = -0.104(10) + 4.718.$$

$$\ln(150 - 68) = -k(3) + 4.718 \quad \ln(T - 68) = 3.678$$

$$4.407 \approx -3k + 4.718 \quad T - 68 \approx 39.6$$

$$k \approx 0.104 \quad T \approx 108$$

PTS: 6 REF: 011139a2 STA: A2.A.28 TOP: Logarithmic Equations

KEY: advanced

428 ANS:

101.43, 12. $r^2 = 25^2 + 85^2 - 2(25)(85)\cos 125.$

$$r^2 \approx 10287.7$$


$$r \approx 101.43$$

$$\frac{2.5}{\sin x} = \frac{101.43}{\sin 125}$$

$$x \approx 12$$

PTS: 6 REF: fall0939a2 STA: A2.A.73 TOP: Vectors

429 ANS:



$$R = \sqrt{28^2 + 40^2 - 2(28)(40)\cos 115} \approx 58 \quad \frac{58}{\sin 115} = \frac{40}{\sin x}$$

$$x \approx 39$$

PTS: 6

REF: 061439a2

STA: A2.A.73

TOP: Vectors