ADVANCED ALGEBRA

Wednesday, January 28, 1948 - 9.15 a. m. to 12.15 p. m., only

Part I

Answer all questions on this part. Each correct answer will receive 24 credits. No partial credit will be allowed. Each answer must be reduced to its simplest form.

1. Write the equation of the straight line passing through the point (4, 3) and parallel to the line $2y = 5x - 2$ 2. In how many points does the graph of $y = x^2 + 3$ intersect the graph of $xy = 12$? 3. Find, to the nearest hundredth, the value of $\sqrt[3]{.456}$ 4. Solve for x the equation $3^x = \frac{1}{6}$ 5. Express $\frac{3+2i}{2-i}$ in the form $a+bi$. 6. The complex numbers $2+2i$ and $1+i\sqrt{3}$ are represented graphically by points A and B . If lines are drawn from A and B to the origin O , find the number of degrees in angle AOB . 7. Write in simplest form the fifth term of $(x+\frac{2}{-})^8$ 8. Find the remainder when $x^{26}-5$ is divided by $x+1$ 9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and $\sqrt{5}$ 10. Transform the equation $x^3-5x^2-7x+2=0$ into an equation whose roots are the roots of the given equation each multiplied by 3. 11. Transform the equation $x^3+5x^2-20x-10=0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3-3x^2+2=0$? 13. Two of the roots of the equation $x^3-5x+2=0$ 0 are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 2 1 centimeters, find the number of seconds in the period when the length is 2 2 centimeters. 15. Find the value of $2x^0+(\frac{x}{x})^2+\frac{x}{x^2}$ when $x=27$ 16. If $f(x)=3x^3-2x+5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be selected from a group of 12 men?		
3. Find, to the nearest hundredth, the value of $\sqrt[4]{.456}$ 4. Solve for x the equation $3^x = \frac{1}{16}$ 5. Express $\frac{3+2i}{2-i}$ in the form $a+bi$. 5. Express $\frac{3+2i}{2-i}$ in the form $a+bi$. 6. The complex numbers $2+2i$ and $1+i\sqrt{3}$ are represented graphically by points A and B . If lines are drawn from A and B to the origin O , find the number of degrees in angle AOB . 7. Write in simplest form the fifth term of $(x+\frac{2}{-})^8$ 8. Find the remainder when $x^{26}-5$ is divided by $x+1$ 9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and $\sqrt{5}$ 10. Transform the equation $x^3-5x^2-7x+2=0$ into an equation whose roots are the roots of the given equation each multiplied by 3. 11. Transform the equation $x^3+5x^2-20x-10=0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3-3x^2+2=0$? 13. Two of the roots of the equation $x^3-px+q=0$ are 4 and 3 . Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 2 centimeters, find the number of seconds in the period when the length is 2 centimeters. 15. Find the value of $2x^0+(\frac{1}{2})^2+\sqrt[3]{x^2}$ when $x=27$ 16. If $f(x)=3x^3-2x+5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be	the point (4, 3) and parallel to the line $2y = 5x - 2$ 2. In how many points does the graph of $y = x^2 + 3$	1
4. Solve for x the equation $3x = \frac{1}{16}$ 5. Express $\frac{3+2i}{2-i}$ in the form $a+bi$. 6. The complex numbers $2+2i$ and $1+i\sqrt{3}$ are represented graphically by points A and B. If lines are drawn from A and B to the origin O, find the number of degrees in angle AOB. 7. Write in simplest form the fifth term of $(x + \frac{2}{x})^8$ 8. Find the remainder when $x^{26} - 5$ is divided by $x + 1$ 9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and √5 10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into an equation whose roots are the roots of the given equation each multiplied by 3. 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation exact diminished by 1. 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{y}{2})^2 + \sqrt[3]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is −96. Find the fourth term. 18. How many different committees of 4 men each can be	· - ·	
5. Express $\frac{3+2i}{2-i}$ in the form $a+bi$. 6. The complex numbers $2+2i$ and $1+i\sqrt{3}$ are represented graphically by points A and B . If lines are drawn from A and B to the origin O , find the number of degrees in angle AOB . 7. Write in simplest form the fifth term of $(x+\frac{2}{x})^8$ 8. Find the remainder when $x^{26}-5$ is divided by $x+1$ 9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and $\sqrt{5}$ 10. Transform the equation $x^3-5x^2-7x+2=0$ into an equation whose roots are the roots of the given equation each multiplied by 3 . 11. Transform the equation $x^3+5x^2-20x-10=0$ into an equation whose roots are those of the original equation each diminished by 1 . 12. How many negative roots has the equation $x^3-3x^2+2=0$? 13. Two of the roots of the equation $x^3-px+q=0$ are 4 and 3 . Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 2 centimeters, find the number of seconds in the period when the length is 2 centimeters. 15. Find the value of $2x^0+(\frac{1}{x})^2+\sqrt[3]{x^2}$ when $x=27$ 16. If $f(x)=3x^3-2x+5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be		-
5. Express — in the form $a + bi$. 6. The complex numbers $2 + 2i$ and $1 + i\sqrt{3}$ are represented graphically by points A and B . If lines are drawn from A and B to the origin O , find the number of degrees in angle AOB . 7. Write in simplest form the fifth term of $(x + \frac{2}{-})^8$ 8. Find the remainder when $x^{26} - 5$ is divided by $x + 1$ 9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and $\sqrt{5}$ 10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into an equation whose roots are the roots of the given equation each multiplied by 3 . 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1 . 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3 . Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 2 centimeters, find the number of seconds in the period when the length is 2 centimeters. 15. Find the value of $2x^0 + (\frac{9}{-})^{-2} + \sqrt[3]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be		-
sented graphically by points A and B . If lines are drawn from A and B to the origin O , find the number of degrees in angle AOB . 7. Write in simplest form the fifth term of $(x + \frac{2}{-})^8$ 8. Find the remainder when $x^{26} - 5$ is divided by $x + 1$ 9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and $\sqrt{5}$ 10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into an equation whose roots are the roots of the given equation each multiplied by 3 . 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1 . 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3 . Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 81 centimeters, find the number of seconds in the period when the length is 81 centimeters, find the number of seconds in the period when the length is 81 centimeters find the number of seconds in the period when the length is 81 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^2 + \sqrt[4]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be	5. Express $\frac{3+2i}{2-i}$ in the form $a+bi$.	5
7. Write in simplest form the fifth term of $(x + \frac{2}{x})^8$ 7. 8. Find the remainder when $x^{26} - 5$ is divided by $x + 1$ 9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and $\sqrt{5}$ 9. 10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into an equation whose roots are the roots of the given equation each multiplied by 3. 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[q]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be	sented graphically by points A and B . If lines are drawn from A and B to the origin O , find the number of degrees in	6
8. Find the remainder when $x^{28} - 5$ is divided by $x + 1$ 9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and $\sqrt{5}$ 10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into an equation whose roots are the roots of the given equation each multiplied by 3. 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[3]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be		0
8. Find the remainder when $x^{28} - 5$ is divided by $x + 1$ 9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and $\sqrt{5}$ 10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into an equation whose roots are the roots of the given equation each multiplied by 3. 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[3]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be	7. Write in simplest form the <i>fifth</i> term of $(x +)^8$	7
9. Write the equation of lowest possible degree with rational coefficients two of whose roots are 2 and $\sqrt{5}$ 10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into an equation whose roots are the roots of the given equation each multiplied by 3. 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^2 + \sqrt[3]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be		
coefficients two of whose roots are 2 and $\sqrt{5}$ 10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into an equation whose roots are the roots of the given equation each multiplied by 3. 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[3]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be		8
10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into an equation whose roots are the roots of the given equation each multiplied by 3. 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[3]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be		9
each multiplied by 3. 11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[q]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be	10. Transform the equation $x^3 - 5x^2 - 7x + 2 = 0$ into	/
11. Transform the equation $x^3 + 5x^2 - 20x - 10 = 0$ into an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[q]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be		10
an equation whose roots are those of the original equation each diminished by 1. 12. How many negative roots has the equation $x^3 - 3x^2 + 2 = 0$? 13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{9}{x})^2 + \sqrt[8]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be		10
12. How many negative roots has the equation x³ - 3x² + 2 = 0? 13. Two of the roots of the equation x³ - px + q = 0 are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of 2x⁰ + (-x/2) -2 + √√x² when x = 27 16. If f(x) = 3x³ - 2x + 5, find f(2). 17. The third term of a geometric series is 12 and the sixth term is -96. Find the fourth term. 18. How many different committees of 4 men each can be	an equation whose roots are those of the original equation	
13. Two of the roots of the equation $x^3 - px + q = 0$ are 14 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of $2x^0 + (\frac{y}{x^2})^{-2} + \sqrt[q]{x^2}$ when $x = 27$ 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be	each diminished by 1.	11
13. Two of the roots of the equation $x^3 - px + q = 0$ are 4 and 3. Find the third root. 14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of 2x ⁰ + (y - 2 + √√x ² when x = 27 16. If f(x) = 3x ³ - 2x + 5, find f(2). 17. The third term of a geometric series is 12 and the sixth term is -96. Find the fourth term. 18. How many different committees of 4 men each can be	12. How many negative roots has the equation $r^3 - 3r^2 + 2 = 0$?	12
14. The period of a pendulum varies directly as the square root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 15. Find the value of 2x ⁰ + (-y/x) ⁻² + √√x ² when x = 27 16. If f(x) = 3x ³ - 2x + 5, find f(2). 17. The third term of a geometric series is 12 and the sixth term is -96. Find the fourth term. 18. How many different committees of 4 men each can be	13. Two of the roots of the equation $x^3 - px + q = 0$ are	
root of its length. If the period is 2 seconds when the length is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 14		13
is 81 centimeters, find the number of seconds in the period when the length is 25 centimeters. 14. 15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[8]{x^2}$ when $x = 27$ 15. 16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be		
15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[8]{x^2}$ when $x = 27$ 15		
16. If $f(x) = 3x^3 - 2x + 5$, find $f(2)$. 17. The third term of a geometric series is 12 and the sixth term is -96 . Find the fourth term. 18. How many different committees of 4 men each can be	the length is 25 centimeters.	14
sixth term is —96. Find the <i>fourth</i> term. 17——— 18. How many different committees of 4 men each can be	15. Find the value of $2x^0 + (\frac{9}{x})^{-2} + \sqrt[8]{x^2}$ when $x = 27$	15
sixth term is —96. Find the <i>fourth</i> term. 17——— 18. How many different committees of 4 men each can be	16. If $f(r) = 3r3 = 2r + 5$ find $f(2)$	16
sixth term is —96. Find the <i>fourth</i> term. 17——— 18. How many different committees of 4 men each can be	17. The third term of a geometric series is 12 and the	+ V
	sixth term is —96. Find the fourth term.	17
selected from a kroup of 15 men.		10
	selected from a group of 12 ment	10

19. How many different numbers each ending in 3 and greater than 1000 can be written with the digits 1, 2, 3 and 4 if no digit is repeated in any number?

20. In a group of children there are 4 boys and 3 girls. If

20. In a group of children there are 4 boys and 3 girls. If cane child is selected by lot, what is the probability that a girl will be selected?

20

Part II

Answer five questions from part II.

21. Find, to the *nearest tenth*, the positive root of $2r^3-6x-1=0$ [10]

22. Solve completely the equation $x^4 + x^5 - x^2 - 7x - 6 = 0$ [10]

23. The number n of vibrations per second made by a stretched string is

given by the relation $n = \frac{1}{2L} \sqrt{\frac{Mg}{m}}$ where L is the length of the string,

M the weight used to stretch the string, m the weight of one centimeter of the string and g = 980. Using logarithms, find n to the nearest integer, when M = 7285 grams, L = 76.8 centimeters and m = 0.0059 gram. [10]

24. a. Using the same set of axes, draw the graphs of $(x + 1)^2 + y^2 = 9$ and $x^2 - y^2 = 4$ [4, 4]

b. From the graphs made in answer to a, find the negative value of x and the negative value of y that satisfy both equations. [Express any approximate value to the nearest tenth.] [2]

25. A number of young men purchased a camp, each paying the same amount. If there had been two more men in the company, each would have said \$12 less. If there had been three fewer men in the company, each would have paid \$24 more. How many men were there and how much did each pay? [10]

26. Three men, A, B and C, set out from the same place and traveled ever the same route at the rates of 4, 5 and 6 miles per hour, respectively. B started 2 hours later than A. How long after B started must C have set out if B and C overtook A at the same moment? [10]

27. Prove that if a + bi is a root of $x^3 + px + q = 0$, in which p and q are real numbers, then a - bi is also a root. [The statement that this is a special case of a more general theorem will not be accepted as **proof**.] [10]

*28. a. Express $2(\cos 150^{\circ} + i \sin 150^{\circ})$ in the form a + bi. [4]

b. Express $1 + i\sqrt{3}$ in polar form. [3]

c. Add graphically -6 + 2i and 3 + 3i. [3]

*29. An open box is to be made from a sheet of tin 12 inches square by cutting equal squares from the four corners and bending up the sides.

a. If x represents a side of the square to be cut out and if y represents the volume of the box, express y as a function of x. [2]

b. Find the derivative of y with respect to x. [4]

c. Find the value of x for which the volume of the hox will be a maximum. [4]

^{*}This question is based on one of the optional topics in the syllabus.