10\_\_\_\_

11\_\_\_\_\_

12

## June 22, 1967

## Part I

Answer all questions in this part. Each correct answer will receive 2½ credits. No partial credit will be allowed. Write your answers in the spaces provided.

- 1. When  $x^{25} 3$  is divided by x + 2, there is an integral remainder. If this remainder is a positive integer, write "positive"; if it is a negative integer, write "negative."
  - 2. Express the product of 3 + 2i and 2i in the form a + bi. 2
  - 3. Solve for a in terms of b and c:  $\frac{1}{a} = \frac{1}{b} + \frac{1}{c}$  3\_\_\_\_\_
- 4. How many different groups of three students each may a teacher choose from among 8 students in her class?
- 5. Write the expression  $[4(\cos 30^{\circ} + i \sin 30^{\circ})]^{2}$  in the form  $r(\cos \theta + i \sin \theta)$ .
- 6. Express the repeating decimal 0.1212..., in which the digits 1 and 2 are repeated endlessly, as a rational number.
- 7. Find the value(s) of x which will satisfy the equation  $\sqrt{2x} + 4 = x$ .
- 8. Find the positive value of b such that the graph of  $x^2 + bx + 4 = y$  touches the x-axis in one and only one point.
- 9. Find the fourth term of the geometric progression whose first two terms are  $\frac{1}{\sqrt{a}}$  and 1, respectively.
  - 10. Find all the values of x which satisfy the inequality

$$\frac{7}{6} + \frac{x}{2} < x - \frac{1}{3}$$
.

- 11. How many cars would a dealer have to have on hand in order to show a customer all of the models available from a manufacturer who builds cars in 6 body styles, with a choice of 4 engines, and equipped with either a standard or an automatic transmission?
- 12. The graph of y = f(x) crosses the x-axis just once between x = -0.9 and x = -0.8. If f(-0.9) > 0, f(-0.85) > 0, and f(-0.8) < 0, find to the nearest tenth a negative root of f(x) = 0.
- 13. Write an equation for the axis of symmetry of the graph of  $y = 10x 2x^2$ .
  - 14. Find the rational fractional root of  $6x^3 + x^2 + 5x 2 = 0$ . 14\_\_\_\_\_

15. What is the <i>least</i> possible degree of a rational integral equation with rational coefficients, if 3 of its roots are $-3$ , $2 + 3i$ , and $4 - \sqrt{7}$ ?	
16. What is the sum of the roots of the equation $3x^3 - 12x^2 + x = 0$ ?	16
17. If $y = x^4 - 2x^2 - 4$ , find the average rate of change of y with respect to x as x varies from $x = -1$ to $x = 0$ .	17
18. The tens digit of a two-digit number is 4 less than the units digit. If the units digit is represented by $u$ , express the number in terms of $u$ .	18
Directions (19-24): Indicate the correct completion for each oring by writing the number 1, 2, 3, or 4 in the space provided.	f the follow-
19. Which defines a rational integral function of $x$ ?  (1) $x^{\frac{1}{4}} - 1$ (2) $x^{2} - 1$ (3) $x^{-\frac{1}{4}} - 1$ (4) $x^{-2} - 1$	19
20. The graphs of $y = ax + b$ and $y = cx + d$ are distinct parallel lines if (1) $a \neq c$ and $b \neq d$ (2) $a \neq c$ and $b = d$ (3) $a = c$ and $b \neq d$ (4) $a = c$ and $b = d$	20
21. Given $\log_9 3^* = 8$ . The value of x that satisfies this equation is (1) 16 (2) 2 (3) 3 (4) 4	21
22. When drawn on the same set of axes, the points of intersection of the graphs of $x^2 + 4y^2 = 16$ and $xy = 1$ are located in quadrants (1) I and III (2) I and IV (3) II and III (4) II and IV	22
23. Given that s varies inversely as the square of t. If a value of t is halved, then the corresponding value of s is multiplied by $(1)$ $\frac{1}{2}$ $(2)$ $(2)$ $(3)$ $\frac{1}{4}$ $(4)$ $(4)$	23
24. How many real roots does $x^4 + 6x^2 + 3 = 0$ have? (1) 1 (2) 2 (3) 0 (4) 4	24
Part II	
Answer sixteen questions from this part, 25-48. Each correct ans ceive 2½ credits. No partial credit will be allowed. Questions m based upon optional topics in the syllabus Write your answers in the vided.	arke <b>d *</b> are
25. When drawn on the same set of axes, the graphs of $y = x^3 - x$ and $y = 2x + k$ are tangent to each other at a point whose abscissa is 1. Find the value of $k$ .	25
26. Express the complex number $\sqrt{2} + i\sqrt{2}$ in the polar form $r(\cos \theta + i \sin \theta)$ .	26
27. A root of $x^3 + x^2 - 24 = 0$ lies between 2 and 3. Find this root to the nearest integer.	27
28. In the equation $x^3 - kx^2 - 5x + 10 = 0$ , k is a rational number. If two roots of this equation are $\sqrt{5}$ and 2, find the numerical value of k.	28

29. The illumination $I$ , received by a body from a source of light of strength $S$ , varies directly as $S$ and inversely as the square of the distance $d$ between the light source and the body. If $k$ represents the constant of variation, express $I$ in terms of $S$ , $d$ and $k$ .	29
30. If $f(x) = 3x^3 - 2x^2 + 1$ , find $f(2) - f(3^{-1})$ .	30
31. In the complex number, plane points $P$ and $Q$ represent the complex numbers $i$ and $-1 + i$ , respectively, and $O$ represents the origin. How many degrees are there in angle $POQ$ ?	31
*32. For what value of $k$ will the value of the determinant $\begin{vmatrix} 4 & k & 3 \\ 2 & 0 & 5 \\ -1 & 0 & 6 \end{vmatrix}$ be 17?	*32
33. Write the term of the expansion of $(a + b)^6$ in which the exponent of $a$ is twice the exponent of $b$ .	33
34. An object is moving along a straight line. Its distance $s$ , in feet from a fixed point after $t$ seconds, is given by the equation $s = 5t^2 + t^3$ . Find the value of $t$ when the acceleration is 19 feet per second per second.	34
35. A school committee of 2 members is to be formed by drawing 2 names from a box containing the names of 5 seniors, 4 juniors, and 3 sophomores. What is the probability that the committee will consist of 2 seniors?	35
36. Write an equation of the straight line having an $x$ -intercept of 2 and a $y$ -intercept of $-4$ .	36
37. If the number $\frac{2+3i}{1-3i}$ is expressed in the form $a+bi$	
where $a$ and $b$ are real numbers, what is the value of $a$ ?	37
38. Find the coordinates of the point of inflection of the curve whose equation is $y = x^3 - 12x + 20$ .	38
39. The lengths of 'he sides of a right triangle form an arithmetic progression whose common difference is 3. Find the length of the shorter leg.	39
40. If $y = \log_{10}3$ , find the value of $10^{27}$ .	40
41. Point $P$ in the first quadrant has the coordinates $(2, k)$ . The line segment between $P$ and the origin and the line segment between $P$ and the point $(10, 0)$ are perpendicular. Find the value of $k$ .	41
*42. Transform the equation $r^2 = 2r \sin \theta - 4r \cos \theta$ from polar to rectangular coordinates.	42

Directions (43-48) For each of those chosen, write in the space provided the number preceding the expression that best completes the statement.

- 43. If x < 0, then  $\sqrt{x^2}$  is equal to (1) an imaginary number (2) x(3) a negative number 43 44. If Al is a years old and Bob is b years old, how old was Al when Bob was c years old? (1) a - b + c (2) a - b - c(3) a + b - c(4) b - a + c45. If  $\log x = 3 + \log 3$ , then x equals (1) 33 (2) 6(3) 3,000 (4) 1.003 46. The circle whose equation is  $(x + 1)^2 + (y + 3)^2 = 9$ is tangent to (1) the y-axis only (2) the x-axis only (3) both axes (4) neither axis 47. If the graphs of  $y = 3^x$  and  $y = 3^{-x}$  are drawn on the same
- 47. If the graphs of  $y = 3^x$  and  $y = 3^{-x}$  are drawn on the same set of axes, the graphs will intersect at a point which is on the line whose equation is (1) x = 0 (2) y = 0 (3) y = x (4) y = -x
- 48. The solution of the inequality  $x^2 x 6 > 0$  is (1) -2 < x < 3 (2) -3 < x < 2(3) x < -2 or x > 3 (4) x < -3 or x > 2