



## F.IF.C.7: Graphing Piecewise-Defined Functions

1 Which graph represents  $f(x) = \begin{cases} |x| & x < 1 \\ \sqrt{x} & x \ge 1 \end{cases}$ ?










2 A function is graphed on the set of axes below.



Which function is related to the graph?

1) 
$$f(x) = \begin{cases} x^2, & x < 1 \\ x - 2, & x > 1 \end{cases}$$

which function is related to
$$f(x) = \begin{cases} x^2, x < 1 \\ x - 2, x > 1 \end{cases}$$

$$2) \quad f(x) = \begin{cases} x^2, x < 1 \\ \frac{1}{2}x + \frac{1}{2}, x > 1 \end{cases}$$

3) 
$$f(x) = \begin{cases} x^2, & x < 1 \\ 2x - 7, & x > 1 \end{cases}$$

3) 
$$f(x) = \begin{cases} x^2, x < 1 \\ 2x - 7, x > 1 \end{cases}$$
4) 
$$f(x) = \begin{cases} x^2, x < 1 \\ \frac{3}{2}x - \frac{9}{2}, x > 1 \end{cases}$$

3 When the function  $g(x) = \begin{cases} 5x, x \le 3 \\ x^2 + 4, x > 3 \end{cases}$  is graphed


correctly, how should the points be drawn on the graph for an *x*-value of 3?

- open circles at (3,15) and (3,13)
- closed circles at (3,15) and (3,13)
- an open circle at (3,15) and a closed circle at (3,13)
- a closed circle at (3, 15) and an open circle at (3,13)

F.IF.C.7: Graphing Piecewise-Defined Functions www.jmap.org


4 On the set of axes below, graph the piecewise function:

$$f(x) = \begin{cases} -\frac{1}{2}x, & x < 2\\ x, & x \ge 2 \end{cases}$$




5 Graph the following function on the set of axes below.

$$f(x) = \begin{cases} |x|, & -3 \le x < 1\\ 4, & 1 \le x \le 8 \end{cases}$$




6 Graph the following piecewise function on the set of axes below.

$$f(x) = \begin{cases} |x|, & -5 \le x < 2\\ -2x + 10, & 2 \le x \le 6 \end{cases}$$




7 Graph the function:  $h(x) = \begin{cases} 2x - 3, & x < 0 \\ x^2 - 4x - 5, & 0 \le x \le 5 \end{cases}$ 



8 The function g is defined as

$$g(x) = \begin{cases} |x+3|, & x < -2\\ x^2 + 1, & -2 \le x \le 2 \end{cases}$$


On the set of axes below, graph g(x).



9 At an office supply store, if a customer purchases fewer than 10 pencils, the cost of each pencil is \$1.75. If a customer purchases 10 or more pencils, the cost of each pencil is \$1.25. Let c be a function for which c(x) is the cost of purchasing x pencils, where x is a whole number.

$$c(x) = \begin{cases} 1.75x, & \text{if } 0 \le x \le 9\\ 1.25x, & \text{if } x \ge 10 \end{cases}$$

Create a graph of *c* on the axes below.




A customer brings 8 pencils to the cashier. The cashier suggests that the total cost to purchase 10 pencils would be less expensive. State whether the cashier is correct or incorrect. Justify your answer.

## **F.IF.C.7: Graphing Piecewise-Defined Functions Answer Section**


1 ANS: 2 REF: 081516ai 2 ANS: 2 REF: 081422ai 3 ANS: 4 REF: 081815ai

4 ANS:




REF: 061832ai

5 ANS:




REF: 011530ai

6 ANS:




REF: 061927ai

7 ANS:




REF: 081932ai

8 ANS:



REF: 012332ai

9 ANS:



cashier is correct.

REF: fall1312ai