F.LE.A.4: Logarithmic Equations 2

- 1 If $\log_2 x = -3$, then x is equal to
 - 1) 9 2) -6 3) $\frac{1}{8}$ 4) -8
- 2 What is the solution of the equation $2\log_4(5x) = 3$?
 - 1) 6.4 2) 2.56 3) $\frac{9}{5}$ 4) $\frac{8}{5}$
- 3 If $\log_2 a = \log_3 a$, what is the value of a?
 - 1) 1 2) 2 3) 3 4) 4
- 4 If $\log_8 x = \frac{2}{3}$, find the value of x.
- 5 Solve for *x*: $\log_8(x+1) = \frac{2}{3}$
- 6 Find x if $\log_9 x = \frac{3}{2}$.
- 7 Solve algebraically for *x*: $\log_{27}(2x-1) = \frac{4}{3}$
- 8 If $\log_x 9 = -2$, what is the value of x?
 - 1) 81 2) $\frac{1}{81}$ 3) 3 4) $\frac{1}{3}$
- 9 Which expression could be used to determine the value of y in the equation $log_x 8 = y$?
 - 1) $\frac{\log 8}{x}$ 2) $\frac{\log 8}{\log x}$ 3) $\frac{8}{\log x}$ 4) $\frac{\log x}{\log 8}$

- 10 If $\log_x 5 = \frac{1}{2}$, find the value of x.
- 11 Solve algebraically for *x*: $\log_{5x-1} 4 = \frac{1}{3}$
- 12 If $\log_x \frac{1}{4} = -1$, find *x*.
- 13 If $\log_x \frac{1}{4} = -2$, find *x*.
- 14 If $\log_4 x = 2.5$ and $\log_y 125 = -\frac{3}{2}$, find the numerical value of $\frac{x}{y}$, in simplest form.
- 15 The temperature, T, of a given cup of hot chocolate after it has been cooling for t minutes can best be modeled by the function below, where T_0 is the temperature of the room and k is a constant. $\ln(T T_0) = -kt + 4.718$

A cup of hot chocolate is placed in a room that has a temperature of 68° . After 3 minutes, the temperature of the hot chocolate is 150° . Compute the value of k to the nearest thousandth. [Only an algebraic solution can receive full credit.] Using this value of k, find the temperature, T, of this cup of hot chocolate if it has been sitting in this room for a total of 10 minutes. Express your answer to the *nearest degree*. [Only an algebraic solution can receive full credit.]

F.LE.A.4: Logarithmic Equations 2 Answer Section

1 ANS: 3

REF: 088624siii

2 ANS: 4

$$2\log_4(5x) = 3$$

$$\log_4(5x) = \frac{3}{2}$$

$$5x = 4^{\frac{3}{2}}$$

$$5x = 8$$

$$x = \frac{8}{5}$$

REF: fall0921a2

3 ANS: 1

$$2^x = a$$

Let $\log_2 \alpha = \log_3 \alpha = x$. Then $2^x = \alpha$ and $3^x = \alpha$. Then $2^x = 3^x$, which is true only if x = 0. $2^0 = \alpha$ or

$$3^x = a$$

$$3^0 = \alpha$$

$$1 = \alpha$$

REF: 080720b

4 ANS:

4

REF: 068404siii

5 ANS:

$$\log_8(x+1) = \frac{2}{3}$$

3.

$$x+1=8^{\frac{2}{3}}$$

$$x + 1 = 4$$

$$x = 3$$

REF: 060925b

6 ANS: 27

REF: 018407siii

7 ANS:

$$2x - 1 = 27^{\frac{4}{3}}$$

$$2x - 1 = 81$$

$$2x = 82$$

$$x = 41$$

REF: 061329a2

8 ANS: 4
$$\log_x 9 = -2$$

$$x^{-2} = 9$$

$$x^2 = \frac{1}{9}$$

$$x = \frac{1}{3}$$

REF: 010819b

9 ANS: 2

$$\log_x 8 = y$$

$$x^{y} = 8$$

$$\log x^{y} = \log 8$$

$$y \log x = \log 8$$

$$y = \frac{\log 8}{\log x}$$

REF: 081621a2

10 ANS:

25

REF: 068813siii

11 ANS:

$$(5x - 1)^{\frac{1}{3}} = 4$$

$$5x - 1 = 64$$

$$5x = 65$$

$$x = 13$$

REF: 061433a2

12 ANS:

REF: 089006siii

13 ANS: 2

REF: 019818siii

14 ANS:

800.
$$x = 4^{2.5} = 32$$
. $y^{-\frac{3}{2}} = 125$. $\frac{x}{y} = \frac{32}{\frac{1}{25}} = 800$

$$y = 125^{-\frac{2}{3}} = \frac{1}{25}$$

REF: 011237a2

15 ANS:

$$ln(T-T_0) = -kt + 4.718$$
 . $ln(T-68) = -0.104(10) + 4.718$.

$$\ln(150 - 68) = -k(3) + 4.718 \quad \ln(T - 68) = 3.678$$

$$4.407 \approx -3k + 4.718$$
 $T - 68 \approx 39.6$

$$k \approx 0.104$$
 $T \approx 108$

REF: 011139a2